This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.





IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of: Martin M. Matzuk et al.

Application No.: 09/830,810

Art Unit: 1653

Filed: October 28, 1999

Examiner: Desai, Anandu U

For: OVARY-SPECIFIC GENES AND PROTEINS

1.132 DECLARATION OF DR. MARTIN M. MATZUK

- 1. I, Martin M. Matzuk, do hereby declare and state the following:
- I am a professor in the Department of Pathology at Baylor College of Medicine.
 I am skilled in the fields of fertility and embryonic development. (See attached Curriculum Vitae).
- I am one of the inventors, and I have read the above-captioned patent application, as well as all Examiner's Actions and responses.
- 4. In the present application, myself and the Pei Wang isolated a polynucleotide sequence, SEQ.ID.NO.1. that was specifically and only expressed in the oocytes. This sequence is known as O1-180 and zygote arrest 1 (Zar1). Since the expression of this protein was similar to GDF-9, we assumed that it would have a similar function.
- 5. We performed several experiments to determine the function of Zar1. Such experiments included producing transgenic mice lacking Zar1 or knockout mice (Zar1^{-/-}). The Zar1^{-/-} females were infertile compare to Zar1^{+/-} and Zar1^{+/-} females. (See page 188, second column of Wu et al., Nature Genetics 33:187-191, 2003).
- I assert that the asserted utility in the patent application is a credible utility based upon our recent evidence as indicated in Wu et al., Nature Genetics 33:187-191, 2003.

7. I hereby declare that all statements made herein on my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like are punishable by fine or imprisonment, or both, under Title 18 § 1001 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

7-19-04	Warten Ne Maty
Date	Martin M. Matzuk, M.D., Pl

CURRICULUM VITAE

NAME: ADDRESS: Martin Matthew Matzuk, M.D., Ph.D.

Department of Pathology

One Baylor Plaza Houston, TX 77030

TELEPHONE:

(713) 798-6451 (713) 798-5833

FAX: E-MAIL:

mmatzuk@bcm.tmc.edu

DATE OF BIRTH: PLACE OF BIRTH: January 30, 1960

New Brunswick, New Jersey

EDUCATION:

University of Chicago, B.A. with Honors, 1982 (Biology)

Washington University, M.D., Ph.D., 1989

Thesis: Structure-function studies of the glycoprotein hormones using mutagenesis,

chimeric genes, and gene-transfer

(Advisor, Dr. Irving Boime; Department of Pharmacology)

PROFESSIONAL EXPERIENCE:

Undergraduate Research Assistant, University of Chicago, Dept. of Biochemistry (Advisor, Dr. Nicholas R. Cozzarelli), 1980-1982

Staff Research Associate, University of California, Berkeley, Dept. of Molecular Biology (Advisor, Dr. Nicholas R. Cozzarelli), 1982-1983

Teaching Assistant, Washington University School of Medicine, Pharmacology, 1985-1986

Clinical Pathology Resident, Dept. of Pathology and Laboratory Medicine, Hospital of the University of Pennsylvania, 1990-1991

Pathology Resident and Postdoctoral Fellow, Dept. of Path. and Molecular Genetics, Baylor College of Medicine, 1991-1993

Clinical Instructor, Dept. of Pathology, Baylor College of Medicine, 1992-1993

Director of Clin. Chem. and Diagnostic Immunol., Ben Taub Gen. Hosp., 1993-present

Assistant Professor, Departments of Pathology, Molecular and Human Genetics, and Molecular and Cellular Biology, Baylor College of Medicine, 1993-1995

Associate Professor, Departments of Pathology, Molecular and Human Genetics, and Molecular and Cellular Biology, Baylor College of Medicine, 1995-1998

Professor, Departments of Pathology, Molecular and Human Genetics, and Molecular and Cellular Biology, Baylor College of Medicine, 1998-present

Stuart A. Wallace Chair in Pathology, Baylor College of Medicine, 1999-present

HONORS:

Edmondson Research Fellowship, University of Chicago, 1981

Medical Scientist Training Program, Washington University, 1983-1989

Spencer T. and Ann W. Olin Fellowship, Washington University, 1989

The Dr. Philip Needleman Prize For Excellence in Pharmacology Research;

Washington University, 1990

Resident Award at the 5th Annual Scientific Symposium of Alumni and Friends, Baylor College of Medicine, 1992

John R. Rainey, MD, Resident Award at the 72nd Annual Meeting of the Texas Society of Pathologists, 1993

Experimental Pathologist-in-Training Award, American Society for Investigative Pathology, 1993

The Michael E. DeBakey, M.D., Excellence in Research Award,

Baylor College of Medicine, 1995

The Richard E. Weitzman Memorial Award, The Endocrine Society, 1996

Inaugural Ernst Knobil Lecture, University of Pittsburgh, 1998

The Bruce Stewart Memorial Lecture, American Society for

Reproductive Medicine, 1998

The HypoCCS Award, Eli Lilly, 1999

O'Neal/Spjut Award, Baylor College of Medicine, 2001

NIH MERIT Award, RO1HD33438, May 1, 2001

Pfizer Outstanding Investigator Award, American Society for Investigative Pathology, 2002

The Society for the Study of Reproduction Research Award, 2002

TEACHING AND ADMINISTRATIVE RESPONSIBILITIES:

Co-director, Medical Scientist Training Program, 1995-present

Member, Medical Scientist Training Program Operating Committee, 1993-present

Faculty Member, Developmental Biology Program, 1994-present

Member, Animal Research Committee, 1994-present

Instructor, Mammalian Molecular Genetics, 1993-1996

Instructor, Developmental Biology I, 1994-1996

Instructor, Reproductive Biology, 1997-present

Laboratory Instructor, Medical School Pathology, 1994-1995

Instructor, Pathology Resident Didactic Lecture Series, 1998-present

Advisor, Graduate and Medical Students (Nine), 1994-present

Member, Graduate Student Thesis Committees (Thirty-six), 1993-present

Member of the Medical Staff, Ben Taub General Hospital, 1993-present

Member of the Medical Staff, The Methodist Hospital, 1994-present

Member, BaylorMedCare, 1997-1999

Member, Faculty Search Committee, Molecular and Cellular Biology, 1995-1996

Member, Faculty Search Committee, Molecular & Human Genetics, 1996-1997

Member, Faculty Search Committee, Pathology, 1996-1997, 1997-1998

Member, Intradepartmental Appointments and Promotions Committee, 1999-present Member, Standing Examination Committee, Molecular and Cellular Biology, 2000-

present

Member, Standing Examination Committee, Molecular and Human Genetics, 2002-present

EXTERNAL THESIS ADVISORY COMMITTEES:

"Thesis defense opponent" Dr. Kirsi Kanaanen Student of Dr. Ilpo Huhtaniemi Turku University, Finland, 1996

"Thesis examiner"
Dr. Shyr-Yeu Lin
Student of Dr. David DeKretser
Monash University, Australia, 2003

AD HOC REVIEWER:

Funding Agencies:

NIH, National Institute of Child Health and Human Development NIH, National Cancer Institute The Lalor Foundation March of Dimes Marsden Fund, The Royal Society of New Zealand Michigan Diabetes Research and Training Center, Pilot/Feasibility Grants Michael Smith Foundation for Health Research MINERVA Cooperative Germany/Israel Research Grants National Health and Medical Research Council, Australia National Science Foundation The New Zealand Neurological Foundation North Carolina Biotechnology Center Research Grants Council of Hong Kong United States Department of Defense United States Department of Agriculture The Wellcome Trust

AD HOC REVIEWER:

Journals:

American Journal of Pathology Archives of Medical Research Biochimica et Biophysica Acta Biotechniques

Biology of Reproduction

Cell

Cytokine

Development Developmental Biology

Developmental Dynamics

The EMBO Journal

Endocrine

Endocrine Reviews

Endocrinology European Journal of Endocrinology

Genes and Development

Genome Research

Genomics

Human Reproduction

Human Reproduction Update

Journal of Andrology

Journal of Cell Science

Journal of Clinical Endocrinology and Metabolism

Journal of Clinical Investigation

Journal of Endocrinology

Journal of Hepatology

Mammalian Ĝenome

Mechanisms of Development Molecular and Cellular Biology

Molecular and Cellular Endocrinology

Molecular Endocrinology

Molecular Human Reproduction

Molecular Reproduction and Development

Nature

Nature Genetics

Nature Medicine

Nature Neuroscience

Nature Reviews Molecular Cell Biology

New England Journal of Medicine

Oncogene

Proceedings of the National Academy of Sciences

Proceedings of the Society for Experimental Biology and Medicine

Reproduction

Science

Trends in Endocrinology and Metabolism

Trends in Genetics

INVITED LECTURES:

- UCLA Symposia on Molecular and Cellular Biology, Molecular Biology of Intracellular Protein Sorting and Organelle Assembly, Workshop, Taos, New Mexico, January 30-February 5, 1987
- Gordon Research Conference, Reproductive Tract Biology, Plymouth, New Hampshire, July 6-10, 1992
- European Science Foundation Scientific Program on Developmental Biology, Workshop on the Transforming Growth Factor β Family, Leuven, Belgium, November 19-21, 1992
- Fifteenth Annual Workshop on Cell and Molecular Techniques, "Techniques in Cellular Signaling and Hormone Action", Houston, Texas, February 28-March 4, 1993
- Southeast Regional Developmental Biology Conference, Athens, Georgia, May 14-16, 1993
- Seventy-Fifth Meeting of the Endocrine Society, Plenary Session, "Gene Deletions and Endocrine Diseases", Las Vegas, Nevada, June 9-12, 1993
- Eighth European Workshop on Molecular and Cellular Endocrinology of the Testis, De Panne, Belgium, March 27-31, 1994
- Fifth International Workshop on Multiple Endocrine Neoplasia, Stockholm, Sweden, June 29-July 2, 1994
- Serono Symposia, Tenth Ovarian Workshop: Frontiers in Ovarian Research, Ann Arbor, Michigan, July 21-23, 1994
- International Workshop on Adrenal Glands, Vascular System, and Hypertension, Hong Kong, April 20-22, 1995
- Third Conference of the International Union of Biochemistry and Molecular Biology, Singapore, April 23-27, 1995
- 24th Annual Reproductive Science Centers Director's Meeting, "Reproductive Developmental Biology", sponsored by the Reproductive Sciences Branch of the Center for Population Research, NICHD, San Francisco, July 14-15, 1995
- Recent Progress in Hormone Research, 51st Conference, Stevenson, Washington, July 29-August 3, 1995
- Serono Symposia, Life Cycle of the Ovarian Follicle, Fort Lauderdale, Florida, November 2-5, 1995

Keystone Symposia on Molecular & Cellular Biology,
"Molecular and Developmental Biology of the Extracellular Matrix",
Keystone Colorado, January 5-11, 1996

Triangle Consortium for Reproductive Biology, "The Development of the Reproductive Systems", <u>Keynote Address</u>, Chapel Hill, North Carolina, January 20, 1996

The American Association for the Advancement of Science Annual Meeting and Science Innovation Exposition, Baltimore, Maryland, February 8-13, 1996

The 25th Anniversary of the National Institutes of Child Health and Human Development Centers' Program, NICHD Molecular Endocrinology Workshop, Houston, Texas, May 21, 1996

Gordon Research Conference, Reproductive Tract Biology, Plymouth, New Hampshire, July 7-12, 1996

29th Annual Meeting of the Society for the Study of Reproduction, London, Ontario, July 27-30, 1996

Reproductive Endocrine Symposium, "Ovarian Messengers", University of Louisville, Louisville, Kentucky, September 27, 1996

3rd International Symposium on Biomedical Diagnostic and Prognostic Indicators, International Society for Preventative Oncology, Nice, France, October 26-28, 1996

Ares-Serono International Symposium on "Inhibin, Activin, and Follistatin... Recent Advances and Future Views", Tokushima, Japan, Nov. 9-10, 1996

HHMI-Sponsored Workshop on "Germ line Development" Chevy Chase , Maryland, December 1-4, 1996

American Society of Andrology, Annual Meeting, Baltimore, Maryland, February 23-25, 1997

"Hormones and Men's Health", Monash Medical Center and Prince Henry's Institute, <u>Keynote Speaker</u>, Melbourne, Australia, March 19-21, 1997

Ovarian Cancer Biology Workshop, National Cancer Institute, Bethesda, Maryland, April 6-8, 1997

VIth International Congress of Andrology, Salzburg, Austria, May 25-29, 1997 INVITED LECTURES (cont): Seventy-Ninth Meeting of the Endocrine Society, Plenary Session, "Genetic Determination of Gonadal Development", Minneapolis, Minnesota, June 11-14, 1997

Bohan Visiting Scientist Lecture, University of Kansas Medical Center Kansas City, Kansas, July 14-15, 1997

American Urological Association Summer Research Conference, "Cell Biology of the Testis: Cell-Cell Interactions", Houston, Texas, August 8-10, 1997

Comitato Promotore Telethon,

Telethon Comatti La Distrofia Muscolare E LE Altre Malattie Genetiche, Bologna, Italy, November 16-18, 1997

"Defining Cytokine Biology through Knockout and Transgenic Mouse Models", The National Cancer Institute. December 11, 1997

Society for Gynecological Investigation, Annual Meeting Course, "How to find a gene and beyond ... ", Atlanta, Georgia, March 11-14, 1998

"Reproduction in the 21st Century", International Symposium Sponsored by the Universidad de la Salle, Bogota, Columbia, March 13, 1998

Sixth Congress of the International Society of Gynecological Endocrinology, Plenary Session, "Molecular Biology in Gynecological Endocrinology", Crans-Montana, Switzerland, March 18-22, 1998

"First International Conference on the Genetic Origins of Premature Ovarian Failure", Sponsored by the Section on Women's Health, Developmental Endocrinology Branch, National Institute of Child Health and Human Development, Washington, D.C., April 2-4, 1998

Genetics Institute Discovery Research Retreat, "Signal Transduction", Ogunquit, Maine, May 20-22, 1998

"Frontiers in Reproduction: Molecular and Cellular Concepts and Applications", Woods Hole, Massachusetts, June 24-27, 1998

5th International Pituitary Congress, Naples, Florida, June 28-30, 1998

INVITED LECTURES (cont):

Society for the Study of Fertility, Annual Meeting,

Plenary Session, "Gonadal Development and Function" Glasgow, Scotland, July 6-8, 1998

XIIth Ovarian Workshop, Plenary Session,
"Follicular Development and Oocyte Maturation"
Houston, Texas, August 5-7, 1998

31st Annual Meeting of the Society for the Study of Reproduction, Techniques Workshop, "Gene Knockouts: Practical Considerations and Expectations", College Station, Texas, August 8, 1998

"The Biology of Menopause", Sponsored by the National Institute on Aging and Serono, Newport Beach, California, September 10-13, 1998

Inaugural Ernst Nobil Lecture, The University of Pittsburgh, Center for Research in Reproductive Physiology, Pittsburgh, Pennsylvania, September 16-17, 1998

37th Annual Meeting of the European Society for Paediatric Endocrinology, Plenary Lecture, Florence, Italy, September 24-27, 1998

Cold Spring Harbor Meeting, Germline Development, Cold Spring Harbor, New York, October 1-3, 1998

16th World Congress on Fertility and Sterility and American Society for Reproductive Medicine, 54th Annual Meeting, "Bruce Stewart Memorial Lecture", San Francisco, California, October 4-8, 1998

4th International Dahlem Symposium, "Cellular Signal Recognition and Transduction", Freie Universitat Berlin, Berlin, Germany, October 8-10, 1998

Annual Meeting of the American Society for Gravitational and Space Biology, Plenary Lecture, Houston, Texas, October 28-31, 1998

Workshop on the Regulation of Reproductive Function, Annual Meeting of the Israel Fertility Association, Jerusalem, Israel, April 12-14, 1999

The Third Annual HypoCCS Symposium, "Regulation of Pituitary Hormone Secretion" Venice, Italy, April 15-17, 1999

1999 North American Inhibin and Activin Congress, Chicago, Illinois, May 21-23, 1999

International Workshop on Early Folliculogenesis and Oocyte Formation, Development and Arrest: Basic and Clinical Aspects, Ares-Serono Foundation, London, England, June 10-11, 1999

32nd Annual Meeting of the Society for the Study of Reproduction, Pullman, Washington, July 31-August 3, 1999

Genetic Regulation of Gametogenesis and Development, The Royal Society of

Edinburgh,

Edinburgh, Scotland, September 23-24, 1999

"Reproductive Sciences 2000: Technology in the Service of Biology", The Annual Meeting of the Society of Gynecologic Investigation, Salt Lake City, Utah, February 23-26, 2000

"The Ovary: from Organogenesis to Function to Failure"
Sponsored by the Section on Women's Health
Developmental Endocrinology Branch
National Institute of Child Health and Human Development,
Washington, D.C., March 30-31, 2000

"Molecular Laboratory Research Course for Clinical Endocrinologists", The University of Hong Kong Hong Kong, May 18-20, 2000

"Molecular Medicine for the Practicing Clinician", Centre of Endocrinology and Diabetes Hong Kong, May 21, 2000

"Embryogenesis Begins During Oogenesis. Follicle Development and Oocyte Fate", Bologna, Italy, May 26-27, 2000

Frontiers in Reproduction Symposium, Oocytes and Human Reproduction, "Egg Futures: 2000 and Beyond" Boston, Massachusetts, June 15-17, 2000

Eighty-Second Meeting of the Endocrine Society, Plenary Session,
"Regulation of Inhibin/Activin Action in Reproduction",
Toronto, Canada, June 21-24, 2000

Gordon Research Conference, Gametogenesis and Embryogenesis, New London, Connecticut, July 2-6, 2000

Ares-Serono Foundation Workshop, "Molecular Genetics of Human Reproduction" Heronissos Beach, Crete, September 15-17, 2000

Round Table Conference, "Alternative Approaches to IVF", Sponsored by N.V. Organon, Lisbon, Portugal, September 22-23, 2000

- "Inhibins, Activins, and Follistatin", Ares-Serono Symposia, Melbourne, Australia, October 26-28, 2000
- 11th Congress of the International Society for Endocrinology, Sydney, Australia, October 29-November 3, 2000
- "Montreal Research Day", Sponsored by the University of Montreal and McGill University, Montreal, Canada, January 17, 2001
- European Society of Human Reproduction and Embryology
 "Mammalian oogenesis and folliculogenesis: in vivo and in vitro approaches",
 Lisbon, Portugal, April 6-7, 2001
- XIIth International Workshop on the Development and Function of the Reproductive Organs, Ares-Sorono Foundation, Jerusalem, Israel, April 30-May 3, 2001
- "First Annual Advances in Contraceptive Help", Sponsored by Wyeth Ayerst Pharmaceuticals Charleston, South Carolina, May 20-22, 2001
 - 17th Annual Meeting of the European Society of Human Reproduction and Embryology, Lausanne, Switzerland, July 1-4, 2001
 - FASEB Conference, "The TGF-β superfamily: signaling and development", Tucson, Arizona, July 7-12, 2001
 - 7th Annual Mouse Developmental Genetics Course, Albert Einstein College of Medicine Bronx, NY, August 26 – September 1, 2001
 - Symposium for the Inauguration of The Weitzmann Women's Health Research Center, "Fertility and Gender-Specific Cancer", Rehovot, Israel, November 11, 2001
 - Ernst Schering Research Foundation Workshop
 "The Future of the Oocyte: Basic and Clinical Aspects"
 Berlin, Germany, January 30 February 1, 2002
 - 21st Joint Meeting of the British Endocrine Societies, Harrogate, United Kingdom, April 8-11, 2002
 - Colloque 2002 de la Société Française de Génétique, "Reproduction et lignée germinale: génétique et pathologies, éthique" Pasteur Institute, Paris, France, April 10-11, 2002

Experimental Biology 2002,

"Translating the genome", American Society for Investigative Pathology New Orleans, Louisiana, April 20-24, 2002

Texas Forum of Female Reproduction, Keynote Speaker, Houston. Texas. May 2-3, 2002

CEJKOVICE 2002, "From Oocyte to Embryonic Stem Cell: A Lesson from Pluripotency", Southern Moravia, Czech Republic, June 6 - 9, 2002

Gordon Research Conference, Gametogenesis and Embryogenesis, New London, Connecticut, June 30-July 4, 2002

35th Annual Meeting of the Society for the Study of Reproduction, Baltimore, Maryland, July 28-31, 2002

5th Copenhagen Workshop on Carcinoma in situ and Cancer of the Testis, Copenhagen, Denmark, August 29-31, 2002

TuBS Symposium on Genetic Engineering of Mice for Biology and Disease Models Mauno Koivisto Center, BioCity Turku, Finland, November 28-29, 2002

Testis Workshop, "Functional Genomics of Male Reproduction" Phoenix, Arizona, March 26-29, 2003

Inaugural speaker, "Billie A. Field Memorial Lectureship", College of Veterinary Medicine, University of Illinois at Urbana – Champaign Urbana, Illinois, April 22, 2003

The Twenty-Second Annual University of Kentucky Symposium in Reproductive Sciences, "Functional Genomics and Mammalian Reproduction" Lexington, Kentucky, May 16-17, 2003

International Conference on the Female Reproductive Tract, Monastery of Seeon, Germany, May 30-June 2, 2003

Ares-Serono Workshop on Human Implantation, "Genomics/Proteomics of the ovary", Madrid, Spain, June 28, 2003

"State of the A.R.T." meeting titled "Communication between oocyte and ovary", Madrid, Spain, June 28-29, 2003

- International Ares-Serono Workshop on "Inhibins, Activins, and Follistatins" Siena, Italy, July 3-4, 2003
- FASEB Conference, "TGF-β Superfamily: Signaling and Development", Tucson, Arizona, July 12-17, 2003
- Eppigenetic contribution to our understanding of the oocyte, The Jackson Laboratory Bar Harbor, Maine, August 2-3, 2003
- 59th Annual Meeting of the American Society for Reproductive Medicine, "Reproductive Medicine and ART: Basic Science and Future Promise" San Antonio, Texas, October 11-15, 2003
- Cytokine-regulated gene expression at the crossroads of innate immunity, inflammation and fertility
 New York University School of Medicine
 New York City, October 17-18, 2003
- Annual meeting, Taiwanese Society for Reproductive Medicine Taipei, Taiwan, December 20-21, 2003
- Reproductive Medicine Conference "Updates in Infertility Treatment" Marco Island, Florida, January 22-24, 2004
- Annual Meeting of the "Hinterzartener Kreis fuer Krebsforschung/Cancer Research" "Growth factors, Repair and Cancer" Cadenabbia, Italy, April 22-25, 2004
- Gordon Research Conference, Gametogenesis and Embryogenesis, New London, Connecticut, June 6-11, 2004
- XIII International Workshop on the Devlopment and Function of the Reproductive Organs Copenhagen, Denmark, June 12-16, 2004
- 37th Annual Meeting of the Society for the Study of Reproduction,
 Minisymposium on "Oocyte-somatic cell interactions in folliculogenesis",
 Vancouver, British Columbia, August 1-4, 2004
- Third International Symposium "Assisted Conception and Reproductive Biology:
 Two Perspectives, One Vision"
 Sponsored by Technobios Procreazione
 "Maternal effect genes during preimplantation development",
 Bologna, Italy, November 18-20, 2004

GRANTS AWARDED (* = Current Support):

National Institutes of Health, National Institute of Child Health and Human Development; K11HD00960;

"Developmental and Functional Roles of Activin and Inhibin"; August 1, 1991-July 31, 1996. I declined years 4-5.

National Institutes of Health, National Institute of Child Health and Human Development; NRSA F32HD07539;

"Developmental and Functional Roles of Activin and Inhibin"; Awarded May 1991; I declined.

The Lalor Foundation; "Developmental and Functional Roles of Activin and Inhibin in Mammalian Reproduction"; June 1, 1991-May 31, 1992; Renewal June 1, 1992-May 31, 1993

*National Institutes of Health, National Institute of Child Health and Human Development; R01HD32067: "Functional Analysis of Activins During Development";

August 1, 1994-July 31, 1997; Renewal December 1, 1998 - May 31, 2004; Renewal June 1, 2004-March 31, 2009

*National Institutes of Health, National Cancer Institute; R01CA60651; "Mouse Models to Study Gonadal Tumor Development"; August 1, 1993-May 31, 1997; Renewal June 1, 1997 - March 31, 2001; Renewal April 1, 2001 - March 31, 2006

*National Institutes of Health, National Institute of Child Health and Human Development; R37 HD33438; "Analysis of Reproductive Function Using Transgenic Mice"; May 1, 1996-April 30, 2001; Renewal May 1, 2001 - April 30, 2006

*National Institutes of Health, National Institute of Child Health and Human Development; U54 HD07495 (PI, Dr. Bert O'Malley); "Center for Reproductive Biology Research Grant"; April 1, 1996-March 31, 1998; Renewal April 1, 1998 - March 31, 2004; Renewal April 1, 2004 - March 31, 2009 (years 32-36)

Genetics Institute, Cambridge Massachussetts; "Functional Analysis of Transforming Growth Factor β Superfamily Members - 1"; October 1, 1996-March 31, 1999; Renewal July 1, 1999 – June 30, 2000

Metamorphix, Baltimore, Maryland; "Functional Analysis of Transforming Growth Factor β Superfamily Members - II"; October 1, 1996-September 30, 1997; Renewal February 1, 1998 - January 31, 1999

National Institutes of Health, National Institute of Child Health and Human Development; RO3 HD37231; "Identification and Analysis of Novel Ovarian Genes"; December 15, 1998 - November 30, 2000

GRANTS AWARDED, continued:

*Wyeth-Ayerst, Women's Health Research Institute;
"Analysis of Putative Female Infertility-Associated Genes";
April 1, 1999 – March 31, 2003; Renewal April 1, 2003 – March 31, 2004

Moran Foundation, Lia Lechago Fund for Cardiovascular Research July 1, 1999 – June 30, 2000

*National Institutes of Health, National Institute of Child Health and Human Development; R01 HD42500; "Regulation of Key Processes in Oocyte Biology" April 1, 2003 – March 31, 2008

ADVISORY/EDITORIAL BOARDS, CHAIR, AND CONSULTANT ACTIVITIES:

Children's Nutrition Research Center Advisory Board, Baylor College of Medicine, Houston, Texas, 1997-present

Research Consultant, Genetics Institute, Cambridge, Massachusetts, 1996-2000

Advisory Committee, The Oregon Regional Primate Research Center, The Oregon Health Sciences University, Portland, Oregon, 1998-present

Research Consultant, Wyeth-Ayerst Research, 1999-present

Advisory Committee, Career Awards in the Biomedical Sciences, Burroughs Wellcome Fund, 1999-present

Editorial Board, Journal of Endocrinology, 2001-present

Editorial Board, Molecular Endocrinology, 2001-present

Committee on New Frontiers in Contraceptive Research, Institute of Medicine/National Academy of Sciences, Board on Health Science Policy, 2003-2004

Co-Chair, Advisory Committee, Career Awards in the Biomedical Sciences, Burroughs Wellcome Fund, 2004-2006

Co-Chair, 3rd Advances in Contraceptive Health Conference, Sponsored by Wyeth Research, Tucson, AZ February 22-24, 2004

ELECTED SOCIETIES:

Sigma Xi, 1982 The Endocrine Society, 1991-present American Society for Investigative Pathology, 1993-present Society for the Study of Reproduction, 1999-present

PATENTS:

Boime, I. and Matzuk, M.M. "Modified Forms of Reproductive Hormones"; U.S. Patent and Trademark Office; Patent Number 5,177,193; Issued January 5, 1993.

Boime, I. and Matzuk, M.M. "DNA encoding reproductive hormones and expression using a minigene"; U.S. Patent and Trademark Office; Patent Number 5,405,945; Issued April 11, 1995.

Boime, I., Matzuk, M.M., and Keene, J.L. "Follicle stimulating hormone-glycosylation analogs", U.S. Patent and Trademark Office, Patent Number 6,306,654; Issued October 23, 2001

Wakil, S., Matzuk, M.M., and Abu-Elheiga, L. "ACC2-knockout mice and uses thereof", U.S. Patent and Trademark Office; Patent Number 6,548,738; Issued April 15, 2003

Matzuk, M.M., Elvin, J.A., and Wang, P. "Assay for growth differentiation factor 9", U.S. Patent and Trademark Office, Patent Number 6,680,174; Issued January 20, 2004

Wakil, S.J., Matzuk, M.M., and Abu-Elheiga, L. "Acetyl-coenzyme a carboxylase 2 as a target in the regulation of fat burning, fat accumulation, energy homeostasis, and insulin action", U.S. Patent and Trademark Office; Patent Number 6,734,337; Issued May 11, 2004

PUBLICATIONS:

- Dean, F., Krasnow, M.A., Otter, R., Matzuk, M.M., Spengler, S.J., and Cozzarelli, N.R. Escherichia coli type-1 topoisomerases: identification, mechanism, and role in recombination. 1. Cold Spring Harbor Symposia on Quantitative Biology, Vol. 47, 769-777 (1983).
- Krasnow, M.A., Matzuk, M.M., Dungan, J.M., Benjamin, H.W., and Cozzarelli, N.R. Sitespecific recombination by Tn3 resolvase: models for pairing of recombination sites. In Mechanisms of DNA Replication and Recombination, (N.R. Cozzarelli, ed.), pp. 637-659, A.R. 2. Liss, New York (1983).
- Benjamin, H.W., Matzuk, M.M., Krasnow, M.A., and Cozzarelli, N.R. Recombination site selection by Tn3 resolvase: topological tests of a tracking mechanism. Cell 40, 147-158 (1985). 3.
- Matzuk, M.M. and Saper, C.B. Preservation of hypothalamic dopaminergic neurons in Parkinson's disease. Ann. Neurol. 18, 552-555 (1985). 4.
- Corless, C.L., Matzuk, M.M., Ramabhadran, T.V., Krichevsky, A., and Boime, I. Gonadotropin beta subunits determine the rate of assembly and the oligosaccharide processing of hormone 5. dimer in transfected cells. J. Cell Biol. 104, 1173-1181 (1987).
- Matzuk, M.M., Krieger, M., Corless, C.L., and Boime, I. Effects of preventing O-linked glycosylation on the secretion of human chorionic gonadotropin in Chinese hamster ovary cells. 6. Proc. Natl. Acad. Sci. USA 84, 6354-6358 (1987).
- Matzuk, M.M., Kornmeier, C.M., Whifield, G.K., Kourides, I.A., and Boime, I. The glycoprotein α subunit is critical for secretion and stability of the thyrotropin β subunit. Molec. 7. Endocrinol. 2, 95-100 (1988).
- **Matzuk, M.M.** and Boime, I. The role of the asparagine-linked oligosaccharides of the α subunit in the secretion and assembly of human chorionic gonadotropin. J. Cell Biol. 106, 1049-1059 8. (1988).
- Campbell, R.K., Matzuk, M.M., Canfield, R.E., Boime, I., and Moyle, W.E. Use of monoclonal antibodies and mutagenesis to study the structure of human chorionic gonadotropin. In Placental 9. Protein Hormones (M. Mochizuki and R. Hussa, eds.), pp. 123-132, Elsevier Science Publishers B.V., Amsterdam (1988).
- Matzuk, M.M. and Boime, I. Site-specific mutagenesis defines the intracellular role of the asparagine-linked oligosaccharides of chorionic gonadotropin β subunit. J. Biol. Chem. 263, 10. 17106-17111 (1988).
- Matzuk, M.M., Keene, J.L., and Boime, I. Site specificity of the chorionic gonadotropin Nlinked oligosaccharides in signal transduction. J. Biol. Chem. 264, 2409-2414 (1989). 11.
- Keene, J.L., Matzuk, M.M., Otani, T., Fauser, B.C.J.M., Galway, A.W., Hsueh, A.J.W., and Boime, I. Expression of biologically active human follitropin in Chinese hamster ovary cells. J. 12. Biol. Chem. 264, 4769-4775 (1989).
- Matzuk, M.M. and Boime, I. Mutagenesis and gene transfer define site-specific roles of the gonadotropin oligosaccharides. Biol. Reprod. 40, 48-53 (1989). 13.

- Bielinska, M., Matzuk, M.M., and Boime, I. Site-specific processing of the N-linked oligosaccharides of the human chorionic gonadotropin α subunit. <u>J. Biol. Chem.</u> 264, 17113-117118 (1989).
- Matzuk, M.M., Spangler, M.M., Camel, M., Suganuma, N., and Boime, I. Mutagenesis and chimeric genes define determinants in the β subunits of human chorionic gonadotropin and lutropin for secretion and assembly. <u>J. Cell Biol.</u> 109, 1429-1438 (1989).
- Suganuma, N., Matzuk, M.M., and Boime, I. Effects of disulfide bond interruption on the assembly and secretion of human chorionic gonadotropin β subunit. <u>J. Biol. Chem.</u> 264, 19302-19307 (1989).
- Keene, J.L., Matzuk, M.M., and Boime, I. Expression of recombinant human choriogonadotropin in Chinese hamster ovary glycosylation mutants. <u>Molec. Endocrinol.</u> 3, 2011-2017 (1989).
- Matzuk, M.M., Hsueh, A.J.W., LaPolt, P., Tsafriri, A., Keene, J.L., and Boime, I. The biological role of the carboxyl-terminal extension of human chorionic gonadotropin β subunit. <u>Endocrinol</u>. 126, 376-383 (1990).
- Moyle, W.R., Matzuk, M.M., Campbell, R.K., Cogliani, E., Dean-Emig, D.M., Krichevsky, A., Barnett, R.W., and Boime, I. Localization of residues which confer antibody binding specificity using human chorionic gonadotropin/luteinizing hormone β subunit chimeras and mutants. <u>J.</u> <u>Biol. Chem.</u> 265, 8511-8518 (1990).
- 20. Campbell, R.K., Matzuk, M.M., Dean-Emig, D.M., Cogliani, E., Myers-Shamy, R.V., Krichevsky, A., Boime, I., Barnett, R., and Moyle, W.R. Use of β-subunit chimeras to study the structures of glycoprotein hormones and to develop a model of the β-subunit. In Glycoprotein Hormones: Structure, Synthesis, and Biological Function (W.W. Chin and I. Boime, eds.), pp. 37-43, Serono Symposia, USA, Norwell, MA (1990).
- Matzuk, M.M., Bielinska, M., and Boime, I. Mutagenesis and gene transfer define the subunitspecific role of the gonadotropin asparagine-linked oligosaccharides. In <u>Glycoprotein Hormones:</u> <u>Structure, Synthesis, and Biological Function</u> (W.W. Chin and I. Boime, eds.), pp. 111-121, Serono Symposia, USA, Norwell, MA (1990).
- Matzuk, M.M., Shlomchik, M., and Shaw, L.M. Making digoxin therapeutic drug monitoring more effective. <u>Therapeutic Drug Monitoring</u> 13, 215-219 (1991).
- Matzuk, M.M. and Bradley, A. Cloning of the human activin receptor cDNA reveals high evolutionary conservation. <u>Biochimica et Biophysica Acta</u> 1130, 105-108 (1992).
- Matzuk, M.M. and Bradley, A. Structure of the mouse activin receptor type II gene. <u>Biochem. Biophys. Rsch. Commun.</u> 185, 404-413 (1992).

- Matzuk, M.M., Finegold, M.J., Su, J.-G.J., Hsueh, A.J.W., and Bradley, A. α-Inhibin is a tumour suppressor gene with gonadal specificity in mice. <u>Nature</u> 360, 313-319 (1992).
- Matzuk, M.M. and Bradley, A. Identification and analysis of tumour suppressor genes using transgenic mouse models. <u>Seminars in Cancer Biology</u> 5, 37-45 (1994).
- Trudeau, V.L., Matzuk, M.M., Hache, R.J.G., and Renaud, L.P. Overexpression of activin-βA subunit mRNA is associated with decreased activin type II receptor mRNA levels in the testes of α-inhibin-deficient mice. Biochem. Biophys. Rsch. Commun. 203, 105-112 (1994).
- Keene, J.L., Nishimori, K., Galway, A.B., Matzuk, M.M., Hsueh, A.J.W., and Boime, I. Recombinant deglycosylated human FSH is an antagonist of human FSH action in cultured rat granulosa cells. <u>Endocrinology Journal</u> 2, 175-180 (1994).
- Incerti, B., Dong, J., Borsani, G., and Matzuk, M.M. Structure of the mouse growth/differentiation factor 9 gene. <u>Biochimica et Biophysica Acta</u> 1222, 125-128 (1994).
- Shikone, T., Matzuk, M.M., Perlas, E., Finegold, M.J., Vale, W., Bradley, A., and Hsueh, A.J.W. Characterization of gonadal tumor cell lines from inhibin-α deficient mice: the role of activin as an autocrine growth factor. <u>Molecular Endocrinology</u> 8, 983-995 (1994).
- Vassalli, A., Matzuk, M.M., Gardner, H.A.R., Lee, K.-F., and Jaenisch, R. Activin/inhibin βB subunit gene disruption leads to defects in eyelid development and female reproduction. Genes and Development 8, 414-427 (1994).
- Matzuk, M.M. Transgenic animals and the study of gonadal function. In <u>Molecular and Cellular Endocrinology of the Testis</u> (G. Verhoeven and U.-F. Habenicht, eds.), pp. 251-271, Springer-Verlag, Berlin (1994).
- Matzuk, M.M., Finegold, M.J., Mather, J.P., Krummen, L., Lu, H., and Bradley, A. Development of cancer cachexia-like syndrome and adrenal tumors in inhibin-deficient mice. Proc. Natl. Acad. Sci. USA 91, 8817-8821 (1994).
- Matzuk, M.M. Gene deletion models to understand cytokine function. In <u>Human Cytokines</u>:
 <u>Their Role in Disease and Therapy</u> (B.B. Aggarwal and R.K. Puri, eds.), pp. 651-662, Blackwell Scientific Publications, Inc., Cambridge, MA. (1995).
- Kumar, T.R., Donehower, L.A., Bradley, A., and Matzuk, M.M. Transgenic mouse models for tumour suppressor genes. <u>J. Internal Medicine</u> 238, 233-238 (1995).
- Matzuk, M.M. Functional analysis of mammalian members of the transforming growth factor β superfamily. <u>Trends in Endocrinology and Metabolism</u> 6, 120-127 (1995).
- Matzuk, M.M., Kumar, T. R., Vassalli, A., Bickenbach, J.R., Roop, D.R., Jaenisch, R., and Bradley, A. Functional analysis of activins in mammalian development. <u>Nature</u> 374, 354-356 (1995).
- Matzuk, M.M., Kumar, T. R., and Bradley, A. Different phenotypes for mice deficient in either activins or activin receptor type II. <u>Nature</u> 374, 356-360 (1995).

- Matzuk, M.M., Lu, H., Vogel, H., Sellheyer, K., Roop, D.R., and Bradley, A. Multiple defects and perinatally death in mice deficient in follistatin. <u>Nature</u> 374, 360-363 (1995).
- Feng, W., Matzuk, M.M., Mountjoy, K., Bedows, E., Ruddon, R.W., and Boime, I. The asparagine-linked oligosaccharides of the human chorionic gonadotropin β subunit facilitate correct disulfide bond pairing. <u>J. Biol. Chem.</u> 270, 11851-11859 (1995).
- Matzuk, M.M., Finegold, M.J., Mishina, Y., Bradley, A., and Behringer, R. R. Synergistic effects of inhibins and Mullerian inhibiting substance on testicular tumorigenesis. <u>Molecular Endocrinology</u> 9, 1337-1345 (1995).
- Kumar, T.R., Kelly, M., Mortrud, M., Low, M.J., and Matzuk, M.M. Cloning of the mouse β gonadotropin subunit genes, I. Structure of the follicle-stimulating hormone β-subunitencoding gene. GENE 166, 333-334 (1995).
- Kumar, T.R. and Matzuk, M.M. Cloning of the mouse gonadotropin β subunit genes, II. Structure of the luteinizing hormone β-subunit-encoding gene. GENE 166, 335-336 (1995).
- Matzuk, M.M., Kumar, T.R., Shou, W., Coerver, K.A., Lau, A., Behringer, R.R., and Finegold, M.J. Transgenic models to study the roles of inhibins and activins in reproduction, oncogenesis, and development. <u>Recent Progress in Hormone Research</u> 51, 123-157 (1996).
- Coerver, K.A., Woodruff, T.K., Finegold, M.J., Mather, J.P., Bradley, A., and Matzuk, M.M. Activin signaling through activin receptor type II causes the cachexia-like symptoms in inhibindeficient mice. <u>Molecular Endocrinology</u> 10, 534-543 (1996).
- Lau, A., Nishimori, K., and Matzuk, M.M. Structure of the mouse activin βC gene. <u>Biochimica</u> et <u>Biophysica Acta</u> 1307, 145-148 (1996).
- Nishimori, K. and Matzuk, M.M. Transgenic mice in the analysis of reproductive development and function. <u>Reviews of Reproduction</u> 1, 203-212 (1996).
- Lieberman, M.W., Wiseman, A.L., Shi, Z-Z., Carter, B.Z., Barrios, R., Ou, C-N., Chevez-Barrios, P., Wang, Y., Habib, G.M., Goodman, J.C., Huang, S.L., Lebovitz, R.M., and Matzuk, M.M. Growth retardation and cysteine deficiency in y-glutamyl transpeptidase-deficient mice. <u>Proc. Natl. Acad. Sci. USA</u> 93, 7923-7926 (1996).
- Kumar, T.R., Wang, Y., and Matzuk, M.M. Gonadotropins are essential modifier factors for gonadal tumor development in inhibin-deficient mice. <u>Endocrinology</u> 137, 4210-4216 (1996).
- Lebovitz, R.M., Zhang, H., Vogel, H., Cartwright, Jr., J., Dionne, L., Lu, N., Huang, S., and Matzuk, M.M. Neurodegeneration, myocardial injury, and perinatal death in mitochondrial superoxide dismutase-deficient mice. <u>Proc. Natl. Acad. Sci. USA</u> 93, 9782-9787 (1996).
- Roberts, V.S., Bentley, C.A., Guo, Q., Matzuk, M.M., and Woodruff, T.K. Tissue-specific binding of radiolabeled activin-A by activin receptors and follistatin in postimplantation rat and mouse embryos. <u>Endocrinology</u> 137, 4201-4209 (1996).

- Nishimori, K., Young, L.J., Guo, Q., Wang, Z., Insel, T.R., and Matzuk, M.M. Oxytocin is required for nursing but is not essential for parturition or reproductive behavior. <u>Proc. Natl.</u> <u>Acad. Sci. USA</u> 21, 11699-11704 (1996).
- Mishina, Y., Rey, R., Finegold, M.J., Matzuk, M.M., Josso, N., Cate, R.L., and Behringer, R.R. Genetic analysis of the Müllerian-inhibiting substance signal transduction pathway in mammalian sexual differentiation. Genes and Development 10, 2577-2587 (1996).
- Dong, J., Albertini, D.F., Nishimori, K., Kumar, T.R., Lu, N., and Matzuk, M.M. Growth differentiation factor-9 is required during early ovarian folliculogenesis. <u>Nature</u> 383, 531-535 (1996).
- Nilsson, C., Pettersson, K., Millar, R.P., Coerver, K.A., Matzuk, M.M., and Huhtaniemi, I.T. Worldwide frequency of a common genetic variant of luteinizing hormone: an international collaborative research. <u>Fertility and Sterility</u> 67, 998-1004 (1997).
- Kumar, T.R., Wang, Y., Lu, N., and Matzuk, M.M. Follicle stimulating hormone is required for ovarian follicle maturation but not male fertility. <u>Nature Genetics</u> 15, 201-204 (1997).
- Shou, W., Guo, Q., Dong, J., and Matzuk, M.M. Transgenic Models for Studying Adrenal, Ovarian, and Testicular Cancer. In <u>Adrenal Gland, Vascular System</u>, and <u>Hypertension</u> (G.P. Vinson and D.C. Anderson, eds.), Journal of Endocrinology Ltd., Bristol, England, pp. 281-292 (1997).
- Young, L.J., Winslow, J.T., Wang, Z., Gingrich, B., Guo, Q., Matzuk, M.M., and Insel, T.R. Gene targeting approaches to neuroendocrinology: oxytocin, maternal behavior and affiliation. <u>Hormones and Behavior</u> 31, 221-231 (1997).
- 59. Lau, A.L., Shou, W., Guo, Q., and Matzuk, M.M. Transgenic approaches to study the functions of the transforming growth factor β superfamily members. In <u>Inhibin. Activin. and Follistatin</u> (T. Aono, H. Sugino, and W.W. Vale, eds.), Springer, New York, pp. 220-243 (1997).
- Shou, W.S., Woodruff, T.K. and Matzuk, M.M. Role of androgens in testicular tumor development in inhibin-deficient mice. <u>Endocrinology</u> 138, 5000-5005 (1997).
- Ben-Arie, N., Bellen, H.J., McCall, A.E., Gordadze, P.R., Armstrong, D.L., Guo, Q., Matzuk, M.M., and Zoghbi, H.Y. Math1 is essential for genesis of cerebellar granule neurons. Nature 390, 169-172 (1997).
- Zhou, J., Kumar, T.R., Matzuk, M.M., and Bondy, C. Insulin-like growth factor I regulates gonadotropin responsiveness in the murine ovary. <u>Molecular Endocrinology</u> 11, 1924-1933 (1997).
- Guo, Q., Kumar, T.R., Woodruff, T.K., Hadsell, L., DeMayo, F., and Matzuk, M.M.
 Overexpression of mouse follistatin causes reproductive defects in transgenic mice. <u>Molecular Endocrinology</u> 12, 96-106 (1998).
- Li, Q., Karam, S.M., Coerver, K.A., Matzuk, M.M., and Gordon, J.I. Stimulation of activin receptor II signaling pathways inhibits differentiation of multiple gastric epithelial cell lineages. Molecular Endocrinology 12, 181-192 (1998).

- Palapattu, G.S., Bao, S., Kumar, T.R., and Matzuk, M.M. Transgenic mouse models for tumor suppressor genes. <u>Cancer Detection and Prevention</u> 22, 75-86 (1998).
- Kumar, T.R. and Matzuk, M.M. Transgenic mice as models of disease. In <u>Textbook of Molecular Medicine</u> (J.L. Jameson, ed.), pp. 97-110, Humana Press, Totowa, New Jersey (1998).
- Draper, L.B., Matzuk, M.M., Roberts, V.J., Cox, E., Weiss, J., Mather, J.P. and Woodruff, T.K. Identification of an inhibin receptor in gonadal tumors from inhibin α-subunit knockout mice. <u>J.</u> <u>Biol. Chem</u> 273, 398-403 (1998).
- Shou, W., Aghdasi, B., Armstrong, D.L., Guo, Q., Bao, S., Charng, M-J., Mathews, L.S., Schneider, M.D., Hamilton, S.L., and Matzuk, M.M. Cardiac defects and altered ryanodine receptor function in mice lacking FKBP12. Nature 391, 489-492 (1998).
- Habib, G.M., Shi, Z.-Z., Cuevas, A., Guo, Q., Matzuk, M.M., and Lieberman, M.W. Leukotriene D₄ and cystinyl-bis-glycine metabolism in membrane-bound-dipeptidase deficient mice. <u>Proc. Natl. Acad. Sci. USA</u> 95, 4859-4863 (1998).
- Ferguson, C.A., Tucker, A.S., Christensen, L., Lau, A.L., Matzuk, M.M., and Sharpe, P.T.
 Activin is an essential early mesenchymal signal in tooth development that is required for
 patterning of the murine dentition. Genes and Development 12, 2636-2649 (1998).
- Matilla, A., Roberson, E.D., Banfi, S., Morales, J., Armstrong, D.L., Burright, E.N., Orr, H.T., Sweatt, J.D., Zoghbi, H.Y. and Matzuk, M.M. Mice lacking ataxin-1 display learning deficits and decreased hippocampal paired-pulse facilitation. <u>Journal of Neuroscience</u> 18, 5508-5516 (1998).
- Kumar, T.R., Low, M.J. and Matzuk, M.M. Genetic rescue of follicle stimulating hormone βdeficient mice. <u>Endocrinology</u> 139, 3289-3295 (1998).
- Jhaveri, S., Erzurumlu, R.S., Chiaia, N., Kumar, T.R., and Matzuk, M.M. Defective whisker follicles and altered brainstem patterns in activin and follistatin knockout mice. <u>Molecular and</u> <u>Cellular Neuroscience</u> 12, 206-219 (1998).
- Bassing, C.H., Shou, W., Muir, S., Heitman, J., Matzuk, M.M., and Wang, X.-F. FKBP12 is not required for the modulation of transforming growth factor beta receptor I signaling activity in embryonic fibroblasts and thymocytes. <u>Cell Growth and Differentiation</u> 9, 223-228 (1998).
- Elvin, J.A. and Matzuk, M.M. Mouse models of ovarian failure. <u>Reviews of Reproduction</u> (1998) 3, 183-195.
- Matzuk, M.M., Dionne, L., Guo, Q., Kumar, T.R., and Lebovitz, R.M. Ovarian function in superoxide dismutase 1 and 2 knockout mice. <u>Endocrinology</u> 139, 4008-4011 (1998).
- Dube, J.L., Wang, P., Elvin, J., Lyons, K.M., Celeste, A.J., and Matzuk, M.M. The bone morphogenetic protein 15 gene is X-linked and expressed in oocytes. <u>Molecular Endocrinology</u> 12, 1809-1817 (1998).

- Carabatsos, M.J., Elvin, J., Matzuk, M.M., and Albertini, D.F. Characterization of oocyte and follicle development in growth differentiation factor-9 deficient mice. <u>Developmental Biology</u> 204, 373-384 (1998).
- Matzuk, M.M. Transgenic models in the study of reproduction. In <u>Fertility and Reproductive</u> <u>Medicine</u> (R.D. Kempers, J. Cohen, A.F. Haney, and J.B. Younger, ed.), pp. 17-25, Elsevier Science B.V. (1998).
- Winslow, J.T., Young, L.J., Hearn, E., Gingrich, B., Wang, Z., Guo, Q., Matzuk, M.M., and Insel, T.R. Phenotypic expression of an oxytocin peptide null mutation in mice. <u>Advances in Experimental Medicine and Biology</u> 449, 241-243 (1998).
- Liu, D., Matzuk, M.M., Sung, W.K., Guo, Q., Wang, P., and Wolgemuth, D.J. Cyclin A1 is specifically required for meiosis in the male. <u>Nature Genetics</u> 20, 377-388 (1998).
- Chang, H., Huylebroeck, D., Verschueren, K., Guo, Q., Matzuk, M.M., and Zwijsen, A. Smad5 knockout mice die at mid-gestation due to multiple embryonic and extraembryonic defects. Development 126, 1631-1642 (1999).
- Matzuk, M.M. and Kumar, T.R. Mouse models to study pituitary hormone gene expression and function. In <u>Regulation of Pituitary Hormone Secretion</u>, (ed. P. Beck-Peccoz), Bioscientifica Ltd., Bristol, U.K., pp. 197-215 (1999).
- Elvin, J.A., Yan, C., Wang, P., Nishimori, K., and Matzuk, M.M. Molecular characterization of the follicle defects in the growth differentiation factor-9-deficient ovary. <u>Molecular</u> <u>Endocrinology</u>, 13, 1018-1034 (1999).
- Elvin, J.A., Clark, A.T., Wang, P., Wolfman, N.M. and Matzuk, M.M. Paracrine actions of growth differentiation factor-9 in the mammalian ovary. <u>Molecular Endocrinology</u>, 13, 1035-1048 (1999).
- Kumar, T.R., Palapattu, G., Wang, P., Woodruff, T.K., Boime, I., Byrne, M.C., and Matzuk, M.M. Transgenic models to study gonadotropin function: the role of FSH in gonadal growth and tumorigenesis. <u>Molecular Endocrinology</u> 13, 851-865 (1999).
- Gold, B.G., Densmore, V., Shou, W., Matzuk, M.M., and Gordon, H.S. Immunophilin FK506-binding protein 52 (not FK506-binding protein 12) mediates the neurotrophic action of FK506.
 <u>The Journal of Pharmacology and Experimental Therapeutics</u> 289, 1202-1210 (1999).
- Lau, A., and Matzuk, M.M. Transgenic models in the study of reproduction. <u>Gravitational and Space Biology Bulletin</u>, 12, 75-82 (1999).
- Elvin, J.A. and Matzuk, M.M. Transgenic models to study reproduction, oncogenesis, and development. In <u>Biology of Menopause</u> (ed. F.L. Bellino), Springer-Verlag, New York, pp. 225-237 (2000).
- Elvin, J.A., Yan, C., and Matzuk, M.M. Oocyte-expressed TGF-β superfamily members in female fertility. <u>Molecular and Cellular Endocrinology</u> 159, 1-5 (2000).

- Ben-Arie, N., Hassan B.A., Bermingham, N.A., Malicki, D.M., Armstrong, D., Matzuk, M.M., Bellen, H.J., and Zoghbi, H.Y. Functional conservation of atonal and Math1 in the CNS and PNS. <u>Development</u> 127, 1039-1048 (2000).
- Kumar, T.R. and Matzuk, M.M. Gene Knockout Models to Study the Hypothalamus-Pituitary-Gonadal Axis. In Gene Engineering and Molecular Models in Endocrinology (M.A. Shupnik, ed.), The Humana Press, Inc., Totowa, NJ, pp. 167-216 (2000).
- Chang, H., Zwijsen, A., Vogel, H., Huylebroeck, D., and Matzuk, M.M. Smad5 is essential for left-right asymmetry in mice. <u>Developmental Biology</u>, 219, 71-78 (2000).
- Davis, A.A., Matzuk, M.M., and Reh, T.A. Activin A promotes progenitor differentiation into photoreceptors in rodent retina. <u>Molecular and Cellular Neuroscience</u>, 15, 11-21(2000).
- Winslow, J.T., Hearn, E.F., Ferguson, J., Young, L.J., Matzuk, M.M., and Insel, T.R. Infant vocalization, adult aggression and fear behavior of an oxytocin null mutant mouse. <u>Hormones and Behavior</u> 37, 145-155 (2000).
- Matzuk, M.M. Revelations of ovarian follicle biology from gene knockout mice. <u>Molecular and Cellular Endocrinology</u> 163, 61-66 (2000).
- Lin, X., Wei, G., Shi, Z., Dryer, L., Esko, J.D., Wells, D.E., and Matzuk, M.M. Disruption of gastrulation and heparin sulfate biosynthesis in EXT1-deficient mice. <u>Developmental Biology</u>, 15, 299-311 (2000).
- Brown, C.W., Houston-Hawkins, D.E., Woodruff, T.K., and Matzuk, M.M. Insertion of *Inhbb* into the *Inhba* locus rescues the *Inhba*-null phenotype and reveals new activin functions.
 <u>Nature Genetics</u> 25, 453-457 (2000).
- Ferguson, J.N., Young, L.J., Hearn, E.F., Matzuk, M.M., Insel, T.R., and Winslow, J.T. Social amnesia in mice lacking the oxytocin gene. <u>Nature Genetics</u> 25, 284-288 (2000).
- Kirkpatrick, L.L., Matzuk, M.M., Dodds, D.C., and Perin, M.S. Biochemical Interactions of the Neuronal Pentraxins: Neuronal Pentraxin (NP) Binds to Taipoxin and Taipoxin-associated calcium-binding protein 49 via NP1 and NP2. J. Biol. Chem. 275, 17786-17792 (2000).
- Cipriano, S.C., Chen, L., Kumar, T.R., and Matzuk, M.M. Follistatin is a modulator of gonadal tumor progression and the activin-induced wasting syndrome in inhibin-deficient mice. <u>Endocrinology</u> 141, 2319-2327 (2000).
- Lorenzetti, D., Watase, K., Xu, B., Matzuk, M.M., and Zoghbi, H.Y. Repeat instability and motor incoordination in mice with a targeted expanded CAG repeat in the Scal locus. <u>Hum. Mol. Genet.</u> 9, 779-785 (2000).
- Pierson, T.M., Wang, Y., DeMayo, F.J., Matzuk, M.M., Tsai, S.Y., and O'Malley, B.W. Regulable expression of inhibin A in wild-type and inhibin α null mice. <u>Molecular Endocrinology</u>, 14, 1075-1085 (2000).

- 104. Shi, Z.Z., Osei-Frimpong, J., Kala, G., Kala, S.V., Barrios, R.J., Habib, G.M., Lukin, D.J., Danney, C.M., Matzuk, M.M., and Lieberman, M.W. Glutathione synthesis is essential for mouse development but not for cell growth in culture. Proc. Natl. Acad. Sci., USA 97, 5101-5106 (2000).
- 105. Lau, A.L., Kumar, T.R., Nishimori, K., Bonadio, J., and Matzuk, M.M. Activin βC and activin βE genes are not essential for mouse liver growth, differentiation, and regeneration. <u>Molecular</u> and <u>Cellular Biology</u>, 20, 6127-6137 (2000).
- Matzuk, M.M. In search of binding-identification of inhibin receptors (Editorial) <u>Endocrinology</u> 141, 2281-2284 (2000).
- Elvin, J.A., Yan, C., and Matzuk, M.M. Growth differentiation factor-9 stimulates progesterone synthesis in granulosa cells via a prostaglandin E2/EP2 receptor pathway. <u>Proc. Natl. Acad. Sci., USA</u> 97, 10288-10293 (2000).
- 108. Kumar, T.R., Wiseman, A.L., Kala, G., Kala, S.V., Matzuk, M.M. and Lieberman, M.W. Reproductive defects in γ-glutamyl transpeptidase-deficient mice. <u>Endocrinology</u> 141, 4270-4277 (2000).
- 109. Yan, C. and Matzuk, M.M. Transgenic models of ovarian failure. <u>Journal of the Society for Gynecologic Investigation</u> 8, 30-33 (2001).
- Chang, H. and Matzuk, M.M. Smad5 is required for primordial germ cell development. <u>Mechanisms of Development</u> 104, 61-67 (2001).
- Elvin, J.A. and Matzuk, M.M. Control of ovarian function. In: <u>Contemporary Endocrinology:</u> <u>Transgenics in Endocrinology</u> (M.M. Matzuk, C.W. Brown, T.R. Kumar, eds.) Humana Press, Totowa, NJ, pp. 61-89 (2001).
- Contemporary Endocrinology: Transgenics in Endocrinology (M.M. Matzuk, C.W. Brown, T.R. Kumar, eds.) Humana Press, Totowa, NJ (2001).
- 113. Wreford, N.G., Kumar, T.R., Matzuk, M.M., and de Kretser, D.M. Analysis of the testicular phenotype of the follicle stimulating hormone (FSH) β subunit knock-out and the activin type IIA receptor knock-out mice by stereological analysis. <u>Endocrinology</u> 142, 2916-2920 (2001).
- 114. Pierson, T.M. and Matzuk, M.M. Transgenic mouse models to study inhibin and activin. In: <u>Inhibins</u>, <u>Activins</u>, and <u>Follistatin in Human Reproductive Physiology</u> (eds. S. Muttukrishna and W. Ledger), Imperial College Press, London, pp. 279-327 (2001).
- Abu-Elheiga, L., Matzuk, M.M., Abo-Hashema, K.A.H., and Wakil, S.J. Continuous fatty acid oxidation and reduced fat storage in mice lacking acetyl-CoA carboxylase 2. <u>Science</u> 291, 2613-2616 (2001).
- 116. Chang, H., Lau, A.L., and Matzuk, M.M. Studying TGF-β superfamily signaling by knockouts and knockins. <u>Molecular and Cellular Endocrinology</u> 180, 39-46 (2001).

- 117. Yan, C., Wang, P., DeMayo, J., DeMayo, F.J., Elvin, J.A., Carino, C., Prasad, S.V., Skinner, S.S., Dunbar, B.S., Dube, J.L., Celeste, A.J., and Matzuk, M.M. Synergistic roles of bone morphogenetic protein 15 and growth differentiation factor 9 in ovarian function. <u>Molecular Endocrinology</u> 15, 854-866 (2001).
- Cipriano, S.C., Chen, L., Burns, K.H., Koff, A., and Matzuk, M.M. Inhibin and p27 interact to regulate gonadal tumorigenesis. <u>Molecular Endocrinology</u> 15, 985-996 (2001).
- 119. Burns, K.H., Yan, C., Kumar, T.R., and Matzuk, M.M. Analysis of ovarian gene expression in follicle stimulating hormone-β knockout mice. <u>Endocrinology</u> 142, 2742-2751 (2001).
- 120. Kumar, T.R., Varani, S., Wreford, N.G., Telfer, N.M., de Kretser, D.M., and Matzuk, M.M. Male reproductive phenotypes in double mutant mice deficient in FSHβ and activin receptor IIA. Endocrinology 142, 3512-3518 (2001).
- Rajkovic, A., Yan, C., Klysik, M., and Matzuk, M.M. Discovery of germ cell-specific transcripts by expressed sequence tag database analysis. <u>Sterility and Fertility</u> 76, 550-554 (2001).
- 122. Jeffs, B., Meeks, J.J., Ito, M., Martinson, F.A., Matzuk, M.M., Jameson, J.L., and Russell, L.D. Blockage of the rete testis and efferent ductules by ectopic Sertoli and Leydig cells causes infertility in Dax1 deficient male mice. Endocrinology 142, 4486-4495 (2001).
- 123. Shi, Z-Z., Han, B., Habib, G.M., Matzuk, M.M., Lieberman, M.W. Disruption of γ-glutamyl leukotrienase results in disruption of leukotriene D₄ synthesis in vivo an attenuation of the acute inflammatory response. Molecular and Cellular Biology 21, 5389-5395 (2001).
- 124. Matzuk, M.M. Eggs in the balance. Nature Genetics 28, 300-301 (2001).
- Durlinger, A.L., Gruijters, M.J., Kramer, P., Karels, B., Kumar, T.R., Matzuk, M.M., Rose, U.M., de Jong, F.H., Uilenbroek, J.T., Grootegoed, J.A., and Themmen, A.P. Anti-Mullerian hormone attenuates the effects of FSH on follicle development in the mouse ovary. Endocrinology 142, 4891-4899 (2001).
- 126. Yan, C., Pendola, F.L., Jacob, R., Lau, A.L., Eppig, J.J., and Matzuk, M.M. Oosp1 encodes a novel mouse oocyte-secreted protein. Genesis 31, 105-110 (2001).
- Wu, X. and Matzuk, M.M. GDF-9 and BMP-15: Oocyte Organizers. <u>Reviews in Endocrine & Metabolic Disorders</u> 3, 27-32 (2002).
- Suzumori, S., Yan, C., Matzuk, M.M., and Rajkovic, A. Nobox is a homeobox-containing gene preferentially expressed in primordial and growing oocytes. <u>Mechanisms of Development</u> 111, 137-141 (2002).
- Rajkovic, A. and Matzuk, M.M. Functional analysis of oocyte-expressed genes using transgenic models. <u>Molecular and Cellular Endocrinology</u> 112, 173-177 (2002).
- 130. Rajkovic, A., Yan, C., Yan, W., Klysik, M, and Matzuk, M.M. Obox, a family of homeobox genes preferentially expressed in germ cells. <u>Genomics</u> 79, 711-717 (2002).

- Rajkovic, A., Lee, J.H., Yan, C., and Matzuk, M.M. The Ret Finger Protein-Like 4 gene, Rfpl4, encodes a putative E3 ubiquitin-protein ligase expressed in adult germ cells. Mechanisms of Development 187, 5-9 (2002).
- Varani, S. and Matzuk, M.M. Phenotypic effects of knockout of oocyte-specific genes. <u>Ernst Schering Research Foundation Workshop</u> 41, 63-79 (2002).
- Burns, K.H., Owens, G., Fernandez, J.M., Nilson, J.H., and Matzuk, M.M. Characterization of integrin expression in the mouse ovary. <u>Biology of Reproduction</u> 67, 743-751 (2002).
- Varani, S., Elvin, J.A., Yan, C., DeMayo, J., DeMayo, F.J., Horton, H.F., Byrne, M.C., and Matzuk, M.M. Knockout of pentraxin 3, a downstream target of growth differentiation factor-9, causes female subfertility. <u>Molecular Endocrinology</u> 16, 1154-1167 (2002).
- Burns, K.H. and Matzuk, M.M. Genetic models for the study of gonadotropin actions. <u>Endocrinology</u> 143, 2823-2835 (2002).
- 136. Yan, W., Rajkovic, A., Viveiros, M.M., Burns, K.H., Eppig, J.J., and Matzuk, M.M. Identification of Gasz, an evolutionarily conserved gene expressed exclusively in germ cells and encoding a protein with four ankyrin repeats, a sterile-alpha motif, and a basic leucine zipper. Molecular Endocrinology 16, 1168-1184 (2002).
- Harlow, C.R., Davidson, L., Burns, K.H., Yan, C., Matzuk, M.M., and Hillier, S.G. FSH and TGF-β superfamily members regulate granulosa cell connective tissue growth factor gene expression in vitro and in vivo. <u>Endocrinology</u> 143, 3316-3325 (2002).
- Matzuk, M.M., Burns, K.H., Viveiros, M.M., and Eppig, J.J. Intercellular communication in the mammalian ovary: occytes carry the conversation. <u>Science</u> 296, 2178-2180 (2002).
- 139. Yan, W., Burns, K.H., Ma, L., and Matzuk, M.M. Identification of Zfp393, a germ cell-specific gene encoding a novel zinc finger protein. <u>Mechanisms of Development</u> 118, 233-239 (2002).
- 140. Burns, K.H. and Matzuk, M.M. Transgenic models for the study of human reproduction. In <u>Biotechnology of Human Reproduction</u> 2nd ed. (A. Revelli, I. Tur-Kaspa, J.G. Holte, and M. Massobrio, eds), pp. 413-435, The Parthenon Publishing Group, New York (2002).
- Chang, H., Brown, C.W., and Matzuk, M.M. Genetic analysis of the mammalian TGF-β superfamily. <u>Endocrine Reviews</u> 23, 787-823 (2002).
- Matzuk, M.M. and Lamb, D.J. Genetic dissection of mammalian fertility pathways. <u>Nature Medicine</u> 8, S41-S49 (2002).
- 143. Burns, K.H., DeMayo, F.J., Matzuk, MM. Transgenic technology, cloning and germ cell transplantation. In Molecular Biology in Reproductive Medicine: Molecular Cellular and Genetic Fundamentals,, second edition (B.C.J.M. Fauser, ed.), pp. 169-195, The Parthenon Publishing Group, New York (2003).
- 144. Nakamura, M., Matzuk, M.M., Gerstmayer, B., Bosio, A., Lauster, R., Miyachi, Y., Werner, S., and Paus, R. Control of pelage hair follicle development and cycling by complex interactions between follistatin and activin. <u>FASEB Journal</u> 17, 497-499 (2003).

- 145. Wu, X., Viveiros, M., Eppig, J.J., Bai, Y., Fitzpatrick, S., and Matzuk, M.M. Zygote arrest 1 (Zar1) is a novel maternal effect gene critical for the oocyte-to-embryo transition. <u>Nature Genetics</u> 33, 187-191 (2003).
- Burns, K.H. and Matzuk, M.M. Oocyte development and maturation. In: <u>Encyclopedia of Hormones</u> (Henry, H.L. and Norman, A.W., eds.), pp. 66-72, Academic Press, San Diego (2003).
- Lukas-Croisier, C., Lasala, C., Nicaud, J., Bedecarras, P., Kumar, T.R., Dutertre, M., Matzuk, M.M., Picard, J.Y., Josso, N., and Rey, R. Follicle-stimulating hormone increases testicular anti-Mullerian hormone (AMH) production through Sertoli cell proliferation and a nonclassical cyclic adenosine 5'-monophosphate-mediated activation of the AMH gene. Molecular Endocrinology 17, 550-561 (2003).
- Suzumori, N., Burns, K.H., Yan, W., and Matzuk, M.M. RFPL4 interacts with oocyte proteins
 of the ubiquitin-proteasome degradation pathway. <u>Proc. Natl. Acad. Sci., USA</u> 100, 550-555
 (2003).
- 149. Wu, M.H., Rajkovic, A., Burns, K.H., Yan, W., Lin, Y.N., and Matzuk, M.M. Sequence and expression of testis-expressed gene 14 (Tex14): a gene encoding a protein kinase preferentially expressed during spermatogenesis. <u>Mechanisms of Development: Gene Expression Patterns</u> 3, 231-236 (2003).
- Yan, W., Burns, K.H., and Matzuk, M.M. Genetic engineering to study testicular tumorigenesis. <u>Acta Pathologica Microbiologica et Immunologica Scandinavica</u> 111, 174-183 (2003).
- Burns, K.H., Viveiros, M.M., Ren, Y., Wang, P., DeMayo, F., Frail, D.E., Eppig, J.J., and Matzuk, M.M. Roles of NPM2 in chromatin and nucleolar organization in oocytes and embryos. <u>Science</u> 300, 633-636 (2003).
- Matzuk, M.M., Hadsell, L.A., De Mayo, F., and Kumar, T.R. Overexpression of human chorionic gonadotropin causes multiple reproductive defects in transgenic mice. <u>Biology of Reproduction</u> 69, 338-346 (2003).
- 153. Chirala, S.S., Chang, H., Matzuk, M.M., Abu-Elheiga, L., Mao, J., Mahon, K., Finegold, M., and Wakil, S.J. De novo fatty acid synthesis is essential for embryonic development: fatty acid synthase null mutants and most of the heterozygotes die in utero. Proc. Natl. Acad. Sci., USA 100, 6358-6363 (2003).
- 154. Yan, W., Assadi, A.H., Wynshaw-Boris, A., Eichele, G., Matzuk, M.M., and Clark, G.D. Previously uncharacterized roles of platelet-activating factor acetylhydrolase 1b complex in mouse spermatogenesis. Proc. Natl. Acad. Sci., USA 100, 7189-7194 (2003).
- Wu, X., Wang, P., Brown, C.A., Zilinski, C.A., and Matzuk, M.M. Zygote arrest 1 (Zar1) is an evolutionarily conserved gene expressed in vertebrate ovaries. <u>Biology of Reproduction</u> 69, 861-867 (2003).
- Brown, C.W., Li, L., Houston-Hawkins, D.E., and Matzuk, M.M. Activins are critical modulators of growth and survival. <u>Molecular Endocrinology</u> 17, 2404-2417 (2003).

- 157. Bernard D.J., Burns, K.H., Haupt, B., Matzuk, M.M., and Woodruff, T.K. Normal reproductive function in lnhBP/p120-deficient mice. <u>Molecular and Cellular Biology</u> 23, 4882-4891 (2003).
- 158. Burns, K.H., Agno, J.E., Chen, L., Haupt, B., Ogbonna, S., Korach, K.S., and Matzuk, M.M. Sexually dimorphic roles of steroid hormone receptor signaling in gonadal tumorigenesis. Molecular Endocrinology 17, 2039-2052 (2003).
- 159. Burns, K.H., Agno, J.E., Sicinski, P., and Matzuk, M.M. Cyclin D2 and p27 are tissue-specific regulators of tumorigenesis in inhibin α knockout mice. <u>Molecular Endocrinology</u> 17, 2053-2069 (2003).
- 160. Yan, W., Ma, L., Burns, K.H., and Matzuk, M.M. HILS1 is a spermatid-specific linker histone H1-like protein implicated in chromatin remodeling during spermiogenesis. <u>Proc. Natl. Acad. Sci., USA</u> 100, 10546-10551 (2003).
- Burns, K.H., Owens, G.E., Ogbonna, S., Nilson, J.H., Matzuk, M.M. Expression profiling analyses of gonadotropin responses and tumor development in the absence of inhibins. <u>Endocrinology</u> 144, 4492-4507 (2003).
- 162. Kumar, T.R., Agno, J., Janovic, J., Conn, P.M. and Matzuk, M.M. Regulation of FSHβ and GnRH receptor gene expression in activin receptor II knockout male mice. <u>Molecular and</u> Cellular Endocrinology 212, 19-27 (2003).
- Burns, K.H. and Matzuk, M.M. The application of gene ablation and related technologies to the study of ovarian physiology. In: <u>The Ovary</u> (Leung, P.C.K. and Adashi, E.Y., eds.), pp. 411-432, Academic Press, San Diego (2004).
- 164. Roy, A., Yan, W., Burns, K.H., and Matzuk, M.M. Tektin3 encodes an evolutionarily conserved putative testicular microtubule-related protein expressed preferentially in male germ cells. Molecular Reproduction and Development 67, 295-302 (2004).
- 165. Yan, W., Ma, L., Matzuk, M.M. Identification and characterization of Gasz orthologs in pufferfish, zebrafish, and frog. <u>Biology of Reproduction</u> 70, 1619-1625 (2004).
- 166. Yao, H.H.C., Matzuk, M.M., Jorgez, C., Menke, D.B., Page, D.C., Swain, A., and Capel, B. Follistatin Operates Downstream of Wnt4 in Mammalian Ovary Organogenesis. <u>Developmental Dynamics</u> 230, 210-215 (2004).
- Jorgez, C.J., Klysik, M., Jamin, S.P., Behringer, R.R., and Matzuk, M.M. Granulosa cellspecific inactivation of follistatin causes female fertility defects. <u>Molecular Endocrinology</u> 18, 953-967 (2004).
- 168. Pangas, S.A. and Matzuk, M.M. Genetic models for transforming growth factor β superfamily signaling in ovarian follicle development. <u>Molecular and Cellular Endocrinology</u>. In press (2004).
- Tang, W., Ingalls, C.P., Durham, W.J., Snider, J., Reid, M.B., Wu, G., Matzuk, M.M. and Hamilton, S.L. Altered excitation-contraction coupling with skeletal muscle specific FKBP12 deficiency. <u>FASEB Journal</u> In press (2004).

- 170. De La Fuente, R., Viveiros, M.M., Burns, K.H., Adashi, E.Y., Matzuk M.M., and Eppig, J.J. Global silencing of transcription in mammalian oocytes does not require major chromatin remodeling. Manuscript submitted.
- 171. Yan, W., Lang, M., Burns, K.H., and Matzuk, M.M. An oocyte-specific oligoadenylate synthase-like protein, OAS1D, inhibits the interferon/oligoadenylate synthetase/RNase Lmediated cell destruction pathway in mouse oocytes. Manuscript submitted.
- 172. Yan, W., Lang, M., Burns, K.H., and Matzuk, M.M. Haploinsufficiency of kelch-like 10 causes infertility in male mice. <u>Proc. Natl. Acad. Sci., USA</u> 101, 7793-7798 (2004).
- 173. Wu, X., Chen, L., Brown, C.A., Yan, C., and Matzuk, M.M. Interrelationship of Growth Differentiation Factor 9 and Inhibin in Early Folliculogenesis and Ovarian Tumorigenesis in Mice. <u>Molecular Endocrinology</u> 18, 1509-1519 (2004).
- 174. Yan, C., Elvin, J.A., Lin, Y-N., Hadsell, L.A., DeMayo, F.J., and Matzuk, M.M. Regulation of growth differentiation factor 9 expression in oocytes in vivo: A key role of the E-box. Manuscript submitted.
- 175. Wang, X., Jorgez, C.J., Suomalainen, M., Matzuk, M.M., Werner, S., and Thesleff, I.. Modulation of TGFβ signaling by follistatin is required for the morphogenesis of mouse molar teeth. Manucript submitted.
- 176. McKenzie, L.J., Jorgez, C.J., Cisneros, P., Buster, J.E., Amato, P., Matzuk, M.M., and Carson, S.A. Follicular fluid prostanoid PGE2: a possible marker for embryo selection? Manuscript submitted.
- Tominaga, K., Ikeno, Y., Ikeda, T., Hawks, C., Matzuk, M.M., Pereira-Smith, O.M. MRG15 regulates embryonic development and proliferation. Manuscript submitted.
- Tominaga, K., Magee, D.M., Matzuk, M.M., and Pereira-Smith, O.M. PAM14, a novel MRG and Rb associated protein, is not required for development and T-cell function in mice. Molecular and Cellular Biology In press (2004).
- Pangas, S.A., Jorgez, C.J., and Matzuk, M.M. Growth differentiation factor 9 regulates expression of the bone morphogenetic protein antagonist, gremlin. <u>Journal of Biological</u> <u>Chemistry</u> In press (2004).
- Combelles, C.M.H., Carabatsos, M.J., Kumar, T.R., Matzuk, M.M., and Albertini, D.F. Hormonal control of somatic cell oocyte interactions during ovarian follicle development. <u>Molecular Reproduction and Development</u> In press (2004).
- 181. Wang, X-P., Suomalainen, M., Jorgez, C.J., Matzuk, M.M., Werner, S., and Thesles, I. Follistatin inhibits ameloblast differentiation by antagonizing BMP4 signaling and is responsible for the enamel-free area formation in mouse incisors. Manuscript submitted.
- 182. Su, Y-Q., Wu, X., O'Brien, M.J., Pendola, F.L., Denegre, J.N., Matzuk, M.M., and Eppig, J.J. Synergistic roles of BMP15 and GDF9 in the development and function of the oocyte-cumulus cell complex in mice: genetic evidence for an oocyte-granulosa cell regulatory loop. Manuscript submitted.

- 183. Ware, S.M., Chen, C., Houston-Hawkins, D.E., Shen, M.M., Matzuk, M.M., and Brown, C.W. Growth differentiation factor 3 regulates gastrulation and embryonic axis formation. Manuscript submitted.
- 184. Wang, Y., Cabello, O.A., Harrison, W., Largaespada, D.A., Matzuk, M.M., and Overbeek, P.A. Rapid identification of transposition in mice using a coat color marker tagged on the Y chromosome. Manuscript submitted.
- Rajkovic, A., Pangas, S.A., Ballow, D., Suzumori, N., and Matzuk, M.M. Nobox regulates oocyte-specific gene expression in early folliculogenesis. Manuscript submitted.
- 186. Ma, X., Dong, Y., Matzuk, M.M., and Kumar, T.R. Targeted disruption of luteinizing hormone β subunit leads to hypogonadism, defects in gonadal steroidogenesis and infertility. Manuscript submitted.
- 187. McKenzie, L.J., Pangas, S.A., Carson, S.A., Kovanci, E., Cisneros, P., Buster, J.E., Amato, P., and Matzuk, M.M. Human granulosa cell gene expression: A predictor of fertilization and embryo selection in women undergoing in vitro fertilization. Manuscript submitted.
- 188. Pangas, S.A., Yan, W., Matzuk, M.M., and Rajkovic, A. Restricted germ cell expression of a gene encoding a novel mammalian HORMA domain-containing protein. Manuscript submitted.
- 189. Burns, K.H. and Matzuk, M.M. Pre-implantation embryogenesis. In <u>The Physiology of Reproduction</u>, 3rd ed. (Neill, J., ed.), pp. _____, Academic Press, San Diego (2004).

Zygote arrest 1 (*Zar1*) is a novel maternal-effect gene critical for the oocyte-to-embryo transition

Xuemei Wu1, Maria M. Viveiros2,5, John J. Eppig2, Yuchen Bai3, Susan L. Fitzpatrick3 & Martin M. Matzuk1,4

Published online 21 January 2003; doi:10.1038/ng1079

The female gamete (the oocyte) serves the distinct purpose of transmitting the maternal genome and other maternal factors that are critical for post-ovulation events 1-4. Through the identification and characterization of oocyte-specific factors, we are beginning to appreciate the diverse functions of oocytes in ovarian folliculogenesis, fertilization and embryogenesis^{5,6}. To understand these processes further, we identified genes called zygote arrest 1 (Zar1 and ZAR1 in mouse and human, respectively) as novel oocyte-specific genes. These encode proteins of 361 amino acids and 424 amino acids, respectively, which share 59% amino-acid identity and an atypical plant homeo-domain (PHD) motif7. Although Zar1-null (Zar1-/-) mice are viable and grossly normal. Zar1-/- females are infertile. Ovarian development and oogenesis through the early stages of fertilization are evidently unimpaired, but most embryos from Zar1-/- females arrest at the one-cell stage. Distinct pronuclei form and DNA replication initiates, but the maternal and paternal genomes remain separate in arrested zygotes. Fewer than 20% of the embryos derived from Zar1-/- females progress to the twocell stage and show marked reduction in the synthesis of the transcription-requiring complex8, and no embryos develop to the four-cell stage. Thus, Zar1 is the first identified oocyte-specific maternal-effect gene that functions at the oocyte-to-embryo transition and, as such, offers new insights into the Initiation of embryonic development and fertility control in mammals.

Pre-implantation embryo development is dependent on stored maternal factors^{1,2}. During meiosis in both sexes, germ-cell genomes are transcriptionally silenced. In the mouse, broad embryonic genome activation occurs at the late two-cell stage^{2,4}. Hence, the oocyte-to-embryo transition depends on maternal transcripts and proteins that accumulate during oogenesis. Despite their many important functions, few mammalian maternal-effect genes have been identified^{9–12}.

To identify novel oocyte-specific genes, we used subtractive hybridization and cDNA library screening to clone a novel gene called zygote arrest 1 (Zarl.), which shows partial homology to a few egg-expressed expressed-sequence tags (AU023153, BB704019). The full-length Zarl cDNA encodes a protein of 51 amino acids (Fig. 1). RT-PCR (Fig. 2a) and northern-blot analyses (data not shown) identified a 1.4-kb Zarl transfer scansies.


only in the ovary. In situ hybridization analysis showed high levels of ZarI in growing oocytes of ovaries from wild-type mice (Fig. 3a,b) and from mice lacking the gene encoding growth differentiation factor 9 (Gdβ); ref. 13; Fig. 3a-f). Consistent with the expressed-sequence tag findings, we were able to amplify ZarI mRNA from wild-type oocytes and one-cell embryos. ZarI mRNA was much less abundant in two-cell embryos and was absent in embryos from four-cell through blastocyst stages and in ZarI^{-1−} oocytes (Fig. 2b). Thus, ZarI mRNA is specifically synthesized in oocytes.

To clone the Zarl gene, we screened genomic libraries and characterized the recovered clones. Zarl and a related pseudogene (Zarl-pst) contain four exons and map to Chromosome 5. Zarl-pst contains agap of 13 nucleotides in exon 1, which is predicted to result in a frameshift and protein truncation in exon 2. RT-PCR with Zarl-specific primers confirmed that Zarl was ovary-specific, but Zarl-psl could not be amplified, establishing it as a pseudogene.



Fig. 1 Comparison of human ZAR1 and mouse Zar1 amino-acid sequences. Highest shared identity was observed in the C termini of these proteins. Asterisks denote the cysteine residues of an atypical PHD motif as predicted from Seasched?

Department of Pathology, Buylor College of Medicine, One Buylor Plaza, Houston, Texas 77030, USA. ²The Lackson Laboratory, Bar Harbor, Maine, USA. ²Weight Research, College-ville, Pennsylvania, USA, ³Department of Molecular and Collund Biology and of Molecular and Human College of Medicine, Houston, Texas, USA, ³Present address: The Center for Transgenesis and Germ Cell Research, Department of Animal Biology, University of Pennsylvania, Romer Square, Pennsylvania, USA. Gerepondenes Should be addressed to M.M. (4-mail: marturk@hom.me. and.)



To identify the human ZARI ortholog, we carried out BLAST searches and genomic-library screens. ZARI, located on Chromosome 4 in a region of conserved synteny to that on mouse Chromosome 5 containing ZarI, spans 4.1 kb and contains four exons, which share 50%, 68%, 84% and 78% nucleotide identity with ZarI exons 1-4, respectively. No human pseudogene was found.

ZAR1 is transcribed exclusively in ovary and testis (Fig. 2c). ZAR1 mRNA is >1.3 kb and encodes a protein of 424 amino acids. ZAR1 and Zar1 proteins share 59% amino-acid identity (Fig. 1), and the C termini, encoded by exons 2-4, are highly conserved (91% shared identity). Although the public database lacks ZAR1 homologs, an atypical PHD motif was discovered in the conserved C termini of ZAR1 and Zar1. PHD sequences are typically characterized by a C4-H-C3 (C-X2-C-X(9-21)-C-X(2-4)-C-X₍₄₋₅₎-H-X₂-C-X₍₁₂₋₄₆₎-C-X₂-C) zinc-binding amino-acid arrangement7. This motif is conserved in the ZAR1 and Zar1 C termini but is a C₈ pattern (C-X₂-C-X₁₃-C-X₂-C-X₄-C-X₁-C-X₁₇-C-X₂-C). A C-to-H substitution is also present in the disease-associated PHD-containing protein ATRX14. Mutagenesis of the H to C in PHD-containing protein KAP-1 has a minimal effect, suggesting that these amino acids are functionally similar15. PHD domains are found in two major classes of proteins: (i) transcriptional activators, repressors or cofactors and (ii) subunits of complexes that modulate chromatin. Thus, ZAR1 may be a transcriptional regulator, and the conserved ZAR1 C terminus is probably functionally important.

Fig. 2. Expression analysis of tygote arrest 1 In mouse and human tissues. All F-PCR analysis of several mouse tissues showed a 26-7 specific 331-9 band only in owary (ov). B, RT-PCR analysis of wild-type (vvt) and Zarr¹ foull express (ou) and wild-type on-ceell (i-t) frough blastoxysic blast) tage embryos detected 2ar1 mRNA in wild-type ocytes and one-cell and two-cell embryos. of RT-PCR analysis of human tissues amplified a fragment of \$30 b, the Through that of APP were used as internal controls for the mouse and human RT-PCR, respectively, as described³². The function of ZARI in the tests (te) is unknown brain; it, lung; he, heart st, stomach; sp. spiene; il, liver; si, small intestine; ki, thinger, ut uterus; o, colons; pp proteste; pl.) pleente; p. papacreas; mu, muscle.

To determine Zarl function, we generated Zarl-null (Zarl^{mulZuk)} mutan mice (Fig. 4). Intercrossing F1 heterozy-gotes (Fig. 4b) yielded 232 F2 progeny (52 wild-type (22.5%), 119 heterozy-gous (Zarl^{+/+}; 51.5%) and 60 homozy-gous-null (Zarl^{-/-}) mice (26.0%)) from 32 litters. Thus, the mutated allele was transmitted with the expected mendelian frequency of 1:2:1. Northern-blot analysis showed a reduction in abundance of Zarl mRNA in Zarl^{+/-} ovaries and was not able to detect the 1.4-kb Zarl mRNA in Zarl^{+/-} ovaries (Fig. 4c), confirming that the Zarl mInXA iled lew sar ull.

 $Zar1^{l+}$ and $Zar1^{l-}$ male and female mice showed no gross or histological abnormalities. To test the fertility of $Zar1^{l+}$ for $Zar1^{l-}$ C781846/19286/89½ hybrid mice, we mated them for six months. Consistent with ovary-specific expression of Zar1, $Zar1^{l-}$ males were fertile (7.4 \pm 0.4 pups per litter). Mating of 14 $Zar1^{l+}$ females with $Zar1^{l+}$ males resulted in 80 litters (9.95 litters per month per mouse; 7.9 \pm 0.3 pups per litter), similar to wild-type matings (8.4 \pm 0.2 pups per litter; ref. 16). In contrast, breeding of 20 $Zar1^{l-}$ females yielded no offspring over six months. Likewise, ten $Zar1^{l-}$ 12956/59½ nibred females bred over six months were also infertile. Thus, Zar1 is executial for female fertility.

To further define Zarl function, we evaluated expression and subcellular localization of Zarl in occytes and early embryos. Consistent with the RT-PCR data, we identified a 45-kDa Zarl-specific band only in mouse ovary by western-blot analysis (Fig. 5a). Zarl localized predominantly to the cytoplasm of occytes of wild-type (Fig. 5b,2) and Gdp9-covaries (Fig. 5d) and was present from primary through antral follide stages (Fig. 5a). Zarl was distributed diffusely throughout the cytoplasm of fully grown occytes isolated




Fig. 3 Localization of Zar1 in moure ovaries. Expression of Zar1 in wild-type (a,b) and Gdf²⁺ (-cf. ref. 13) ovaries treated with PMSG was analyzed by in situ hybridization with a specific antisense probe. Bright-field (a,ce) and corresponding dark-field (b,d) images of each only rection are presented. Areas of panels c and d are shown at higher magnotion in e and f. Zar1 mRNA was detected at early primary foliate (type a) through antial foliate (type 8) taspes, but not in primordial foliates (type 3). The foliate dissification is based on Pederan and Peters²⁷.





Fig. 4 Targeting strategy of Zar1 and Southern-blot and northern-blot analyses of F2 offspring, a, A targeting vector was constructed by replacing exon 1 (the largest exon, which contains the initiation ATG) and part of intron 1 with a PgkHPRT expression cassette. The MCTtk expression cassette was used for negative selection. Targeted ES cell clones containing a wild-type (wt) allele, a pseudogene allele (Zar1-ps1) and a mutated (mut) allele were confirmed by Southern-blot analysis²² and greed to cost contex containing a wilest-type (viz writer, a procusoper where (xor) party and a mountain the containing a wilest-type (viz writer) and the containing a wilest-type (viz writer) and the containing a wilest were the produced produce detected only the pseudogene in DNA derived from Zar1+ mice. c, Northern-blot analysis of ovarian mRNA from willd-type, Zar1+ and Zar1+ females using the fulllength Zar1 cDNA. On longer exposure, a smaller transcript of unknown relevance was observed in Zar1+ ovaries. The expression level of the 1.4-kb Zar1 mRNA in wild-type mice was approximately twice the levels present in Zar1st ovaries, d, Gapd was used as a control for equivalent loading on the northern blot.


from ZarI+/- mice (Fig. 5f) but was not present in ovaries (Fig. 5e) and oocytes (Fig. 5g) from Zar1--- females. Zar1 was also detected after resumption of meiosis through progression to metaphase II (Fig. 5h,i). Zar1 persisted in one-cell embryos (Fig. 5j) but was markedly less abundant in two-cell embryos (Fig. 5k). Thus, Zar1 could function in growing oocytes through formation of two-cell embryos. The rapid disappearance of Zarl at the two-cell stage, however, suggests a critical role in the oocyte-to-embryo transition.

we examined ovarian histology. Ovaries from Zar1-1- females after insemination (82.4 ± 7.5%). Whereas the first cleavage

(Fig. 5e) were indistinguishable from those of control females. All stages of follicle development and corpora lutea (indicative of ovulation) were evident in ovaries of Zar1-/- females. Moreover, superovulation resulted in similar numbers of oocytes from $Zar1^{+/-}$ (31.6 ± 4.7; n = 8) and $Zar1^{-/-}$ (34.3 ± 4.1; n = 14) females. Most $Zar1^{+/-}$ (74.4 ± 5.5%; n = 156) and $Zar1^{-/-}$ oocytes (62.9 \pm 4.3%; n = 137) resumed meiosis and progressed to metaphase II during a 17-h culture. Like heterozygous controls To determine the cause of the infertility in female Zar1-- mice, (92.9 ± 2.1%), Zar1-- oocytes formed two pronuclei within 8 h

Fig. 5 Zar1 expression. A polyclonal antibody against Zar1 was used for western-blot (a), immunohistochemistry (b-e) immunofluorescence analyses (f-k) to detect Zar1 expression. a, Zar1 was detected only in mouse ovary (ov) and was absent in spleen (sp), lung (lu) and testis (te), unlike actin (positive control) A smaller 35-kDa band was also detected in the ovary and may represent a degra dation product. Similar to Zar1 mRNA, Zar1 protein expression (red staining) began in oocytes of primary and continued follicles through all follicle stages in wild-type ovaries (b,c). Zar1 was also detected in GdfS ovaries (d) but not in Zarfovaries (e). Zar1 (shown in green) was detected predominantly in the cytoplasm of grown, prophasi





I-arrested oocytes from Zar1++ (f) but not Zar1++ mice (g). Zar1 was expressed in wild-type oocytes during the progression from MI (h) to MII (i) and persisted in zygotes at the one-cell stage, collected 6 h after fertilization (j). But Zar1 expression was markedly lower in two-cell-stage embryos (k), with bright staining evident only in polar body remnants (arrow). DNA labeling is shown in red, af, antral follicle; cl, corpus luteum.




Fig. 6 Development of embryos derived from Zar1+ and Zar1+ mice. Adult Zar1++ (a) and Zar1++ (b) females were mated with stud males. Eggs and embryos were collected from the oviduct and uteri on d 3.5 after vaginal plug. Whereas all zygotes (n = 41) from Zar1++ female mice (n = 6) progressed to the blastocyst stage (a), most zygotes (n = 19) from $Zar1^{-1}$ mice (n = 9) remained at the one-cell stage, and many degenerating embryos were detected (b).

Oocytes from Zar1+- and Zar1+- mice (c) were matured and fertilized in vitro. 24 h after fertilization, the arrested zygotes from Zar1-f emales were labeled with antibody against 8-tubulin and propidium jodide to assess microtubule and chromatin configurations, respectively. Decondensed chromatin (red) was evident in maternal and paternal pronuclei. The microtubules are shown in green. In a second experiment, the fertilized zygotes were placed in medium with BrdU 8 h after fertilization and cultured overnight (d). Immunofluores-





ence analysis showed BrdU incorporation in both pronudei (bright orange) of an arrested zygote from a Zar1-f- female, indicative of entry into S phase. e, Early embryos from inseminated Zar1 null eggs underwent embryonic genome activation. TRC synthesis was detected in two-cell-stage embryos from wild-type control (ctri) and Zar1+ females. Culture with α-amanitin (alpha-am, 11 µg ml-1), an inhibitor of RNA polymerase II, prevented TRC synthesis. The TRC experiments were performed as described⁸ using 15 embryos per lane.

occurred in most in vivo (89.3%) and in vitro (86.5 ± 1.4%) fer- Notably, inhibition of embryonic transcription does not pretilized embryos from Zar1+/- females, we observed two-cell embryos (some of which appeared fragmented) in only 19.1 ± 9.1% (in vitro) and 20.8 ± 1.5% (in vivo) of embryos from Zar1-females. 100% of embryos isolated from uteri of Zar1+ffemales developed to blastocysts by day 3.5, and most cultured control embryos progressed to morula and blastocyst stages (82.1 ± 1.5%), but we observed only fragmented onecell embryos and a single two-cell embryo in Zar1-i- females (Fig. 6a,b). Therefore, an arrest at the zygote and two-cell stages accounts for the infertility of Zar1-i- females.

Additional analysis of the embryonic block showed that Zar1 null zygotes progressed through G1 and successfully entered S phase. Both pronuclei had decondensed chromatin and incorporation of 5-bromo-2-deoxyuridine (BrdU; Fig. 6c,d). The microtubule network showed an interphase configuration with no assembled spindle apparatus. In vitro fertilized oocytes were treated with colcemid, which depolymerized microtubules and arrested all zygotes from Zar+/- females at M phase. Only a few Zar1 null zygotes were similarly arrested, but this number corresponded to the small percentage of Zar1 null embryos that entered M phase and became two-cell embryos, indicating that most arrested earlier at the S or G2 stages. Thus, the maternal and paternal genomes remained separate in discrete pronuclei, and the two haploid genomes did not unite, indicating that the fertilization had not been completed.

To further define the block in the Zar1 null embryos, we analyzed the synthesis of a group of proteins called the transcriptionrequiring complex (TRC), markers of embryonic genomic activation8. Although TRC proteins were detected in the limited number of Zar1 null embryos that reached the two-cell stage, their TRC expression levels were 15% of those of the control embryos, suggesting that there are defects in embryonic genome activation (Fig. 6e). The precise mechanisms by which Zar1 directs the oocyte-to-embryo transition have yet to be determined

The decrease in Zar1 expression in normal two-cell embryos correlates with the usual rapid degradation of maternal transcripts after fertilization 17. Developmental processes ongoing at this time include completion of the meiotic-to-mitotic cellcycle transition and activation of the embryonic genome¹⁻⁴.

vent the first mitotic division but arrests embryos beyond the two-cell stage17. Of the other maternal-effect genes9-12, only Mater (maternal antigen that embryos require) is expressed exclusively in oocytes and early pre-implantation embryos. Compared with Zar1 null embryos, Mater null embryos block at the two-cell stage and have higher levels of genome activation. The predominant one-cell-stage arrest of fertilized eggs from Zar1-/- females suggests that this maternal factor functions before embryonic genome activation. Although both Mater (53%) and Zarl (59%) share low identity with their human orthologs 18, these human orthologs probably have conserved functions in early embryogenesis and fertility. Identification of proteins that interact with Zar1 may provide insights into the roles of this maternal protein in regulating the essential transition from oocyte to embryo. This study presents the first identification and characterization of zygote arrest 1, a gene expressed during oocyte development, the function of which is key for the initiation of embryogenesis.

Methods

mRNA expression analysis. We obtained total RNA from mouse tissues or embryos using the RNA STAT-60 method (Leedo Medical Laboratories). We carried out in situ hybridization with a Zarl-specific probe as described 19,20. We carried out RT-PCR analysis of mouse and human cDNAs (primer sequences are available upon request).

Mouse ovary cDNA library and genomic library screening. We used a Zar1 cDNA probe (corresponding to nucleotides 896-1,220 of the fulllength Zar1 cDNA) to screen a wild-type mouse ovary cDNA library. We used the full length 1,259-nt Zar1 cDNA, which contains an open reading frame from nt 28-1,110, to screen 129S6/SvEv mouse genomic λ FixII phage and human genomic DNA libraries (Stratagene). We also used Zar1specific primers to screen a 129X1/SvJ BAC library (Genome Systems). We hybridized filters as described21.

ES-cell manipulation and Southern-blot analysis. We electroporated the linearized Zar1 targeting vector into the HPRT-negative AB2.2 ES-cell line, selected clones in hypoxantine, aminopteridine and thimidine and 1-(2'deoxy-2'fluoro-β-D-arabinofuranosyl)-5-iodouracil and sczeened DNA from the clones by Southern-blot analysis. We found that 5 of 171 (3%) of the ES cell clones analyzed were targeted at the Zar1 locus. ES cell line Zar1-140-G11 was injected into blastocysts to produce chimeric male mice

that were fertile and transmitted the mutated Zarl allele (Zarltml Zuk, herein called Zar1") to F1 progeny as described 22,23

Superovulation, mating and in vitro embryo culture. We injected 25-d-old ZarI* and ZarI* female mice with pregnant mare serum gonadotropin (PMSG; intraperitoneal injection of 5 IU per mouse) and gave them human chorionic gonadotropin (intraperitoneal injection of 5 IU per mouse) 48 h later. Mice were then caged overnight with (C57BL/6) × 129S6/SvEv) F1 stud males. The following morning, we recovered eggs and embryos in M2 medium, counted them and cultured them in vitro up to 4 d in M16 medium (Sigma). Alternatively, adult mutant females were mated to stud males, uteri and oviducts were flushed on d 1.5, 2.5 or 3.5, and embryos were collected and cultured in M16 medium.

In vitro oocyte maturation and fertilization. We injected sexually mature Zar1+1- and Zar1-1- female mice with 5 IU of PMSG to stimulate preovulatory follicle development. We isolated cumulus-enclosed oocyte complexes 48 h later and cultured them for 17 h in minimal essential medium with 5% serum. We subsequently removed the surrounding somatic cells and examined the oocytes to determine the progression of meiosis. Mature MII-stage eggs were inseminated in vitro with sperm from wildtype (C57BL/6] × SJL/J) F1 mice24. We assessed development of zygotes and two-cell-stage embryos 6 and 24 h after fertilization, respectively. We evaluated blastocyst formation on d 5

Production of polyclonal antibodies against Zarl and immunostaining We subcloned a partial mouse Zar1 cDNA (nt 178-1,083 corresponding to amino acids 51-352) into pET23b vector (Novagen) and injected chimeric recombinant Zar1 protein (with a T7 tag at the N terminus and His tag at the C terminus) into goats to produce polyclonal antibodies (CoCalico Biologicals). We carried out western-blot analysis and immunostaining as described²⁵ using primary antibody (diluted at 1:1,000). For western-blot analysis, we used secondary antibody against goat conjugated with horseradish peroxidase (Boehringer Mannheim; diluted at 1:2,000) and ECL Western Blotting Detection Reagents (Amersham Biosciences). For immunostaining, we carried out incubation with secondary antibody and visualization of positive cells using the New Fuchsin kit (BioGenex). We used pre-immune serum in control sections.

Immunofluorescence. We carried out immunofluorescence analysis of oocytes and embryos as described25. We carried out reactions with the goat Zarl antisera (diluted at 1:1,000 in block solution) for 1 h and then exposed samples to 3 µg ml-1 of antibody against rabbit IgG conjugated with fluorescein isothiocyanate (Jackson Immuno Research Laboratories) for 45 min. We labeled DNA with propidium iodide (10 µg ml⁻¹ for 10 min). We evaluated negative controls with pre-immune serum

We evaluated fertilized zygotes that had not undergone the first mitotic division by 24 h after fertilization to determine chromatin and microtubule configurations. The zygotes were fixed, permeabilized and blocked. We carried out all subsequent steps, including rinses, at 37 °C in block solution. We labeled microtubules with antibody against β -tubulin (3.8 μg ml-1 for 1 h) and secondary antibody against mouse conjugated with fluorescein isothiocyanate (1.3 µg mH for 45 min) and labeled DNA with propidium iodide as above. We detected fluorescence using a confocal microscope.

DNA synthesis. Fully-grown oocytes from Zar1+i- and Zar1-i- mice were in vitro matured and fertilized as above. Approximately 8 h after fertilization, we transferred the zygotes that formed male and female pronuclei to medium supplemented with 50 µM BrdU for overnight culture. 24 h after fertilization, embryos were fixed and processed to assess BrdU incorporation26. We detected fluorescence using a confocal microscope.

Statistical analysis. We calculated statistical significance by one-way ANOVA. Data are expressed as mean ± s.e.m

URL. ScanSite27 is available at http://scansite.mit.edu

GenBank accession numbers. Zar1 cDNA, AY191415; Zar1 gene, AY193889; ZARI cDNA, AY191416; ZARI gene, AY193890.

Acknowledgments

We thank M. Berry, R. Burgess, K. Burns, T. Gridley, G. Kopf and T. Kumar for critical review of the manuscript, S. Baker for aid in manuscript formatting and C. Brown, K. Burns, L. Chen, L. Erdos, M. O'Brien, P. Wang and C. Yan for technical assistance. These studies were supported in part by a sponsored research grant from Wyeth Research and US National Institutes of Health grants (to M.M.M. and J.J.E.). X. Wu was supported in part by a post-doctoral fellowship from the Lalor Foundation.

Competing interests statement

The authors declare competing financial interests: see the Nature Genetics website (http://www.nature.com/naturegenetics) for details.

Received 20 August; accepted 9 December 2002

- Latham, K.E. Mechanisms and control of embryonic genome activation in
- mammallan embryos. Int. Rev. Cyrol. 193, 71–124 (1999). Latham, K.E. & Schultz, R.M. Embryonic genome activation. Front. Biosci. 6,
- Schultz, R.M. Regulation of zygotic gene activation in the mouse. Bioessays 15, 531-538 (1993)
- Aoki, F., Worrad, D.M. & Schultz, R.M. Regulation of transcriptional activity during the first and second cell cycles in the pre-implantation mouse embryo. Dev. Biol. 1B1, 295-307 (1997).
- Matzuk, M.M., Burns, K., Viveiros, M.M. & Eppig, J. Intercellular communication in the mammalian ovary: oocytes carry the conversation. Science 295, 2178–2180
 - Matzuk, M.M. & Lamb, D.J. Genetic dissection of mammalian fertility pathways
- Matzui, M.M. del., S41-S49 (2002). Nat. Med. B, S41-S49 (2002). & Stewart, A.F. The PHD finger: implications for Assland, R., Gibson, T.J. & Stewart, A.F. The PHD finger: implications for chromatin-mediated transcriptional regulation. Trends Biochem. Sci. 20, 56-59
 - Conover, J.C., Temeles, G.L., Zimmermann, J.W., Burke, B. & Schultz, R.M. Stagespecific expression of a family of proteins that are major products of zygotic gene activation in the mouse embryo. Dev. Biol. 144, 392–404 (1991).
- activation in the mouse emoryo. Job. Diol. 1996, 332-499 (1971). Tong, Z.B. et al. Mater, a maternal effect gene required for early embryonic development in mice. Nat. Genet. 25, 267–268 (2000). ted by a maternal-effect mutation in Howell, C.Y. et al. Genomic Imprinting disrupt the Dnmt1 gene. Cell 104, B29–83B (2001).
- 11. Prolla, T.A. et al. Tumour susceptibility and spontaneous mutation in mice deficient in Mih1, Pms1 and Pms2 DNA mismatch repair. Nat. Genet. 18, 276-279
- 12. Christians, E., Davis, A.A., Thomas, S.D. & Benjamin, I.J. Maternal effect of Hsf1 on (199B). reproductive success. Nature 407, 693-694 (2000).
 - 13. Dong, J. et al. Growth differentiation factor 9 is required during early ovarian folliculogenesis. Nature 383, \$31–535 (1996).
 - Gibbons, R.J. et al. Mutations in transcriptional regulator ATRX establish the functional significance of a PHD-like domain. Nat. Genet. 17, 146–148 (1997). 15. Capill, A.D., Schultz, D.C., Rauscher, L.F. & Borden, K.L. Solution structure of the
 - ארות, אבער, באינות בער המערה (R. & Borden, K. L. Solution structure of the PHD domain from the KAPA Loopersors structural determinants for PHD, RING and LIM across domains. EMBO J. 20, 155–177 (2001).

 16. Kumar, T.R. et al. Improductive defects in y-glutamyl transpeptidase-deficient mice. Embour. 106, 114, 4270–4277 (2000).
 - 17. Flach, G., Johnson, M.H., Braude, R.R., Taylor, R.A. & Bolton, V.N. The transition from maternal to embryonic control in the two-cell mouse embryo. *EMBO J.* 1,
 - 681-686 (19B2). Tong, Z.B., Bondy, C.A., Zhou, J. & Nelson, L.M. A human homologue of mouse Mater, a maternal-effect gene essential for early embryonic develop
 - Reprod. 17, 903-911 (2002). 19. Albrecht, U., Eichele, G., Helms, J.A. & Lu, H.C. Visualization of gene expression
 - patterns by In situ hybridization in Molecular and Cellular Methods in Developmental Toxicology (ed. Daston, G.P.) 23–48 (CRC Press, Boca Raton, Florida, 1997).
 - Florida, 1997).

 G. Elvin, J.A., Yan, C., Wang, P., Nishimori, K. & Matzuk, M.M. Molecular Characterization of the follide defects in the growth differentiation factor specificient ovary. Mol. Endocrinol. 13, 1018–1034 (1999).

 1. Matzuk, M.M. & Bradley, A. Structure of the mouse activin receptor type II gene.
 - Biochem. Biophys. Res. Commun. 1BS, 404-413 (1992). Matzuk, M.M., Finegold, M.J., Su, J. G.J., Hsueh, A.J.W. & Bradley, A. α-Inhibin is a tumor-suppressor gene with gonadal specificity in mice. Nature 360, 313–319
 - Bradley, A. Production and analysis of chimeric mice-teratocarcinomas and embryonic stem cells: a practical approach. in Production and Analysis of Chimeric Mice (ed. Robinson, E.J.) 113–151 (Oxford, London, 1987).
 - Chimeric Mice (ed. Kobinson, E.J.) 113-131 (Oxford, London, 1987).
 24. Eppig, J.J. Mouse oocyte maturation, fertilization, and pre-implantation development in vitro. in A Comparative Methods Approach to the Study of Oocytes and Embryos (ed. Richter, J.D.) 3-9 (Oxford University Press, Oxford, Dec.)
 - 25. Yan, W. et al. Identification of Gasz, an evolutionarily conserved gene express exclusively in germ cells and encoding a protein with four ankyrin repeats, a ste alpha motif, and a basic leucine zipper. Mol. Endocrinol. 16, 1168–1184 (2002). 26. Ferreira, J. & Carmo-Fonseca, M. Genome replication in early mo
 - follows a defined temporal and spatial order. J. Cell. Sci. 110, 889-897 (1997)
 - colouws a defined temporal and spatial order. J. Cell. 201, 110, 899-897 (1997).

 7 Yaffe, M.S. et al. A motif-based profile scanning approach for genome-wide prediction of signaling pathways. Nat. Biotechnol. 19, 348–353 (2001).

 28. Elvin, J.A., Cark, A.T., Wang, P., Wolfman, N.M. & Matzuk, M.M. Paragcine actions of growth differentiation factor 9 in the mammalian ovary. Mol. Endocrinol. 13, 1035-1048 (1999)
 - Pedersen, T. & Peters, H. Proposal for a classification of occytes and follicles in the mouse ovary. J. Reprod. Fertil. 17, 555–557 (1968).